

Six new species of the *Anopheles leucosphyrus* group, reinterpretation of *An. elegans* and vector implications

M. A. M. SALLUM^{1,2}, E. L. PEYTON¹ and R. C. WILKERSON¹

¹Walter Reed Biosystematics Unit, Smithsonian Institution – Museum Support Center, Suitland, Maryland, U.S.A. and

²Departamento de Epidemiologia, Núcleo de Pesquisas Taxonômica e Sistemática em Entomologia Médica, Faculdade de Saúde Pública, Universidade de São Paulo, Brazil

Abstract. Among Oriental anopheline mosquitoes (Diptera: Culicidae), several major vectors of forest malaria belong to the group of *Anopheles (Cellia) leucosphyrus* Dönitz. We have morphologically examined representative material (>8000 specimens from seven countries) for taxonomic revision of the Leucosphyrus Group. Six new species are here described from adult, pupal and larval stages (with illustrations of immature stages) and formally named as follows: *An. latens* n. sp. (= *An. leucosphyrus* species A of Baimai *et al.*, 1988b), *An. cracens* n. sp., *An. scanloni* n. sp., *An. baimaii* n. sp. (formerly *An. dirus* species B, C, D, respectively), *An. mirans* n. sp. and *An. recens* n. sp. Additionally, *An. elegans* (James) is redescribed and placed in the complex of *An. dirus* Peyton & Harrison (comprising *An. baimaii*, *An. cracens*, *An. dirus*, *An. elegans*, *An. nemophilous* Peyton & Ramalingam, *An. scanloni* and *An. takasagoensis* Morishita) of the Leucosphyrus Subgroup, together with *An. baisasi* Colless and the *An. leucosphyrus* complex (comprising *An. balabacensis* Baisas, *An. introlatus* Baisas, *An. latens* and *An. leucosphyrus*). Hence, the former Elegans Subgroup is renamed the Hackeri Subgroup (comprising *An. hackeri* Edwards, *An. pujutensis* Colless, *An. recens* and *An. sulawesi* Waktoedi).

Distribution data and bionomics of the newly defined species are given, based on new material and published records, with discussion of morphological characters for species distinction and implications for ecology and vector roles of such species. Now these and other members of the Leucosphyrus Group are identifiable, it should be possible to clarify the medical importance and distribution of each species. Those already regarded as vectors of human malaria are: *An. baimaii* [Bangladesh, China (Yunnan), India (Andamans, Assam, Meghalaya, West Bengal), Myanmar, Thailand]; *An. latens* [Borneo (where it also transmits Bancroftian filariasis), peninsular Malaysia, Thailand]; probably *An. cracens* (Sumatra, peninsular Malaysia, Thailand); presumably *An. scanloni* (Thailand); perhaps *An. elegans* (the Western Ghat form of *An. dirus*, restricted to peninsular India); but apparently not *An. recens* (Sumatra) nor *An. mirans* [Sri Lanka and south-west India (Karnataka, Kerala, Tamil Nadu)], which is a natural vector of simian malarias. Together with typical *An. balabacensis*, *An. dirus* and *An. leucosphyrus*, therefore, the Leucosphyrus Group includes about seven important vectors of forest malaria, plus at least a dozen species of no known medical importance, with differential specific distributions collectively spanning > 5000 km from India to the Philippines.

Correspondence: Professor Maria Anice Mureb Sallum, Faculdade de Saúde Pública, Universidade de São Paulo, Avenida Dr Arnaldo 715, São Paulo, SP 01246-904, Brazil. Tel. +55 11 3066 7731; fax: +55 11 3282 1898; E-mail: masallum@usp.br

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 2005		2. REPORT TYPE		3. DATES COVERED 00-00-2005 to 00-00-2005	
4. TITLE AND SUBTITLE Six new species of the Anopheles leucosphyrus group, reinterpretation of An. elegans and vector implications				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Walter Reed Biosystematics Unit, Smithsonian Institution, Washington, DC, 20560				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT see report					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 42	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

Key words. *Anopheles baimaii*, *An. cracens*, *An. dirus*, *An. elegans*, *An. latens*, *An. leucosphyrus*, *An. mirans*, *An. recens*, *An. scanloni*, *Plasmodium falciparum*, *P. cynomolgi*, *P. fragile*, *P. inui*, *P. vivax*, *Wuchereria*, Culicidae, filaria vector, forest malaria vectors, malaria transmission, new species, morphology, mosquito systematics, Oriental Region.

Introduction

Among the anopheline mosquito fauna of the Oriental Region, the most important vectors of forest malaria belong to the *Anopheles leucosphyrus* Group (Reid, 1949, 1968), classified in *Neomyzomyia* Series (Christophers, 1924) of subgenus *Cellia* Theobald of *Anopheles* Meigen (Harbach, 2004). The geographical distribution of the Leucosphyrus Group ranges from the southern islands of the Philippines through most of mainland South-east Asia to the Chinese islands of Hainan and Taiwan in the east and westward to southern India and Sri Lanka (Peyton, 1990). Adult males and females of all species in the Leucosphyrus Group have a distinctive broad white-scaled band covering the apex of the hindtibia and base of hindtarsomere I. Taxonomy of the Leucosphyrus Group was first revised by Colless (1956), who recognized six species (including two subspecies) and five geographical forms of doubtful status, two of which were named as subspecies by Colless (1957): *An. balabacensis introlatus* for the Kepong Form and *An. balabacensis baisasi* for the Luzon Form. Peyton (1990) corroborated Colless' classification and recognized 14 formally named species, six unnamed species and two geographical forms; he also subdivided the Leucosphyrus Group into Elegans, Leucosphyrus and Riparis Subgroups, based on differential length ratios of the adult female proboscis/forefemur and maxillary palpus/proboscis. Because we ascertained that *An. elegans* belongs to the *An. dirus* complex of the Leucosphyrus Subgroup, as explained below, we realign the Elegans Subgroup and propose to rename it the Hackeri Subgroup.

As treated in this paper and its sequel (Sallum *et al.*, 2005), the Leucosphyrus Subgroup consists of two widespread and often sympatric complexes of sibling species, generally known as *An. dirus* s.l.* and *An. leucosphyrus* s.l., plus *An. baisasi* Colless in The Philippines and a poorly known form in Con Son island, Vietnam. The Dirus Complex includes *An. dirus* Peyton & Harrison (*An. dirus* s.s.* or species A), three previously unnamed species (*An. dirus* B, C and D), *An. elegans* (= *An. dirus* E), *An. takasagoensis* Morishita and *An. nemophilous* Peyton & Ramalingam. The Leucosphyrus Complex includes *An. balabacensis* Baisas, *An. introlatus* Colless and *An. leucosphyrus* Dönitz

(*An. leucosphyrus* s.s. = species B of Baimai *et al.*, 1988b) plus one previously unnamed species (*An. leucosphyrus* A). The Riparis Subgroup remains unchanged with three recognized species: *An. cristatus* King & Baisas, *An. macarthuri* Colless and *An. riparis* King & Baisas. The Hackeri Subgroup has three recognized species (*An. hackeri*, *An. pujutensis* and *An. sulawesi*) plus two previously unnamed species, one known as *An. leucosphyrus* Sumatra Form and the other previously misidentified as *An. elegans* (Colless, 1956; Reid, 1968; Peyton, 1990; Harbach, 2004).

Peyton & Harrison (1979) described *An. dirus* based on morphological features of the adult, pupal and larval stages. Later, Peyton & Harrison (1980) removed *An. takasagoensis* from synonymy with *An. balabacensis* based on morphological evidence and hybridization tests between *An. dirus* and *An. takasagoensis*. Other studies of the Dirus and Leucosphyrus Complexes showed that they were groups of apparently isomorphic but reproductively distinct species. Discovery of most members of the Dirus Complex was achieved primarily with crossing studies, cytogenetics (chromosome banding patterns), allozyme data, DNA probes and DNA hybridization (Hii, 1984b; Wibowo *et al.*, 1984; Baimai *et al.*, 1987, 1988a; Panyim *et al.*, 1988; Green *et al.*, 1992; Audtho *et al.*, 1995). Cytogenetic studies on members of the Leucosphyrus Group in Thailand and peninsular Malaysia showed differences in the amount and distribution of heterochromatin in the sex chromosomes, suggesting that *An. dirus* s.l. comprised at least four genetically distinct genetic species: these were designated *An. dirus* A, B, C and D (Baimai *et al.*, 1981, 1984; Hii, 1982; Wibowo *et al.*, 1984). Kanda *et al.* (1981, 1983) studied nine geographical populations of the Leucosphyrus Group collected in Thailand and Malaysia: from morphological characters, hybridization tests and cytogenetics they found various degrees of relatedness between the populations, concluding that *An. balabacensis* and *An. dirus* were not specifically distinct. More recently, however, molecular methods (DNA-based polymerase chain reaction, PCR) have been developed to differentiate members of the Dirus Complex (Audtho *et al.*, 1995; Xu *et al.*, 1998; Walton *et al.*, 1999; Huong *et al.*, 2001; Manguin *et al.*, 2002), confirming the status of species A, B, C and D. Also, two species in the Leucosphyrus Group, *An. leucosphyrus* A and B, were recognized from mitotic chromosome karyotypes and cross-breeding experiments (Baimai *et al.*, 1988b).

Hitherto, three members of the Leucosphyrus Group – *An. balabacensis*, *An. dirus* s.l. and *An. leucosphyrus* s.l. – have been considered to be of epidemiological importance because

**Sensu stricto* (s.s.) means 'in the strict sense', i.e. the exact species. *Sensu lato* (s.l.) means 'in the broad sense', i.e. any or all members of the species complex.

they are highly competent vectors of human malaria parasites in South-east Asia. Moreover, *An. balabacensis* and *An. leucosphyrus* s.l. transmit *Wuchereria* parasites causing lymphatic filariasis in Borneo (Zulueta, 1957; White, 1983). However, the involvement of members of the *Leucosphyrus* Group in the epidemiology of human malaria and filariasis is not fully understood because the species involved in transmission are difficult to distinguish using morphological characters. These species can be identified only by examining morphological characters of associated adult, pupal and larval stages, or by genetic methods (chromosomal characters and molecular diagnostics). To facilitate identification of known species, in preparation for taxonomic revision of the *Leucosphyrus* Group (Sallum *et al.*, 2005), this paper formally describes and names six new species based on morphological features, and settles the taxonomic position of *An. elegans*.

Materials and methods

We employed standard methods for mounting, storing and microscopic examination of anopheline mosquitoes (<http://wrbu.si.edu/wrbu.html>), with emphasis on adults associated with pupal and larval skins, after their collection from the field by many collaborators using various sampling methods (Service, 1993). Terminology for morphological characters follows Harbach & Knight (1980; 1981), except we used Belkin (1962) for wing veins and Wilkerson & Peyton (1990) for wing spots. The toothed margin index of the pupal paddle was defined by Colless (1956) to express the ratio of the lengths: (a) from the tip of the inner basal tubercle of the paddle to the base of the most distant marginal tooth and (b) from the same point to the base of seta 1-Pa. The basal extension of the PSD spot (see Wing spot abbreviations) on vein R as compared to the HD, PSP and PSD spots on the costa defines the level of extension of the PSD (Peyton & Ramalingam, 1988). Level 1 is defined as the PSD on vein R not extending basally beyond the PSD on the costa, level 2 as the PSD extending basally but not beyond 0.5 of the PSP, level 3 as the PSD extending basally beyond 0.5 of the PSP but not to the level of the HD, level 4 as the PSD extending basally to the level of the HD but not beyond 0.5 of the HD, level 5 as the PSD extending basally beyond 0.5 of the HD but not beyond the HD, and level 6 as the PSD extending basally beyond the HD. The number of pale interruptions on PSD-PD spots is defined as the sum of pale interruptions on PSD and PD spots of vein R.

Abbreviations

Acc. no., collection accession number; AFRIMS, Armed Forces Research Institute of Medical Sciences, Bangkok, Thailand; ASA, allele-specific amplification; BMNH, The Natural History Museum, London, U.K.; F, female;

H-banding, heterochromatin bands of the chromosome; ITS2, internal transcribed spacer 2 of the ribosomal DNA; L, larva; Le, larval exuvia; M, male; mtDNA, mitochondrial DNA; NMNH, National Museum of Natural History, Washington D.C., U.S.A.; P, pupa; PCR, polymerase chain reaction; Pe, pupal exuvia; RFLP, restriction fragment length polymorphism; SCAR, sequence characterized amplified regions. Wing spots: AD, apical dark spot; ASP, accessory sector pale spot; HD, humeral dark spot; HP, humeral pale spot; PD, posterior dark spot; PHP, prehumeral pale spot; PP, preapical pale spot; PSD, presector dark spot; PSP, presector pale spot; SCP, subcostal pale spot; SD, sector dark spot; SP, sector pale spot. An asterisk (*) after the life stage means that at least part of it is illustrated. Complete lists of specimens examined and distribution maps for each species will be provided later (Sallum *et al.*, 2005), along with illustrations of relevant parts of adults, fourth-instar larva and pupa, showing taxonomic character states, plus identification keys to the *An. leucosphyrus* group of species.

Leucosphyrus subgroup

Leucosphyrus Complex

Anopheles (Cellia) latens Sallum & Peyton, sp.n.

Etymology: the name *latens* is from Latin, meaning concealed, secret, hidden, latent.

Anopheles leucosphyrus of Leicester, 1903: 291 (bionomics).

Myzomyia leucosphyrus of Daniels, 1908: 1 (*An. hackeri* and *An. leucosphyrus* mixed, bionomics); Leicester, 1908a: 7 (bionomics); Leicester, 1908b: 18 (F, M, bionomics, probably mixed with *An. hackeri*); Leicester, 1908c: 267 (feeding behaviour, probably mixed with *An. hackeri*).

Neomyzomyia elegans of James & Stanton, 1912: 59.

Anopheles leucosphyrus of Kanda *et al.*, 1981: 321 (hybridization, taxonomy); Kanda *et al.*, 1983: 193 (phylogeny, chromosomal polymorphism); Takai *et al.*, 1984: 145 (genetics); Takai, 1986: 45 (*in part*, electrophoresis, Sarawak); Harbach *et al.*, 1987: 241 (bionomics); Chang *et al.*, 1995: 192 (bionomics); Seng, 1999: 454 (bionomics).

Anopheles leucosphyrus species A of Baimai *et al.*, 1988b: 44 (cytogenetics, crossbreeding); Peyton, 1990: 197 (taxonomy).

Female. Head: proboscis uniformly dark-scaled, length 1.92–2.25 mm (mean 2.11 mm), ratio of length to forefemur length 1.06–1.16 (mean 1.12), maxillary palpus length 1.62–2.15 mm (mean 1.89 mm), ratio of length to proboscis length 0.81–0.95 (mean 0.89), ratio of length to forefemur length 0.91–1.07 (mean 1.00), ratio of length of palpomeres 3/4 1.62–2.08 (mean 1.77), 3/5 2.17–2.87 (mean 2.44), 4/5 1.22–1.62 (mean 1.37), 4–5/3 0.84–1.08 (mean 0.98), palpomeres 2 and 3 with narrow apical band of silvery white

scales, apical pale band of palpomere 4 variable, absent, smaller or larger than those of palpomeres 2 and 3, scales pale cream-coloured to golden, slightly contrasting with those on palpomeres 2 and 3, palpomere 5 dark-scaled at base, with apical band of pale cream-coloured to golden scales, length of apical pale band 0.17–4.00 (mean 1.18) length of basal dark band of palpomere 5, ratio of length of apical pale band of palpomere 4 to length of basal dark band of palpomere 5 0.0–1.67 (mean 0.51). Thorax: pleural setae as follow: 1–4 upper proepisternal, 0–2 prespiracular, 5–8 prealar, 4–6 upper mesokatepisternal, 2,3 lower mesokatepisternal, 4–7 upper mesepimeral. Wing: length 3.16–3.62 mm (mean 3.46 mm), pale yellow scales on costa and subcosta, pale scales on remaining veins white, not strongly contrasting with others; PHP spot of costa small, always present, HP spot always present, usually prominent, rarely very small, PSP spot always present and prominent, rarely reduced, SP spot usually small, sometimes prominent, ASP spot usually present, occasionally reduced or absent, PP spot 0.55–1.54 (mean 1.03) length of SCP spot, AD spot 1.17–3.00 (mean 1.84) length of PP spot, PSD spot of vein R extending basally from level 4 to level 6 on one or both wings, with 1–5 small pale interruptions, SD spot of vein R with 1–4 pale interruptions, sum of pale spots on PSD-PD interruptions of vein R 4–12 for each wing, ratio of length of cell R_2 to vein R_{2+3} 1.46–2.00 (mean 1.69), ratio of length of cell R_2 to cell M_{1+2} 1.19–1.34 (mean 1.28). Legs: femora, tibiae and tarsomeres 1 dark-scaled, speckled with spots of pale scales; foretarsomere 2 with apical and basal pale bands, middle region dark-scaled, sometimes with 1–4 pale spots on dorsal surface, foretarsomeres 3 and 4 mostly dark-scaled, with basal and apical pale bands, foretarsomere 4 sometimes mostly pale-scaled with inconspicuous middle band of light brown scales, foretarsomere 5 variable, mostly pale-scaled with inconspicuous patch of tan scales on dorsal surface, or with basal dark band, pale-scaled at apex, dark scales on ventral surface of foretarsomeres 2–5 light brown to tan; midtarsomeres 2–5 mostly dark-scaled with narrow apical bands of pale scales; hindtarsomeres 2–5 dark-scaled with narrow apical bands of pale scales, hindtarsomeres 2, 3 sometimes with few pale spots in middle of dark area. Abdomen: tergum VI frequently without scales, sometimes with 2, 3 golden scales on posterolateral margin, tergum VII with few cream-coloured to golden scales on posterior margin or rarely without scales, tergum VIII covered with yellowish to golden scales; sternum VI without scales, sternum VII with a posteromedial patch of dark scales, sternum VIII with small lateral patch of whitish scales.

Male. Essentially as in female except for sexual characters. Wing generally paler with reduced scaling, pale spots usually longer than in female. Palpomere 2 with patch of pale scales at middle of dorsal surface, extending to lateral surface, apex of palpomere 2 bare; palpomere 3 dark-scaled, with long dorsal patch of pale scales at middle, extending to lateral surface, apex of palpomere 3 with broad band of pale scales interrupted by short patch of dark scales at apex of ventrolateral surface; palpomeres 4 and 5 mostly pale-scaled

with narrow basal band of dark scales. Abdomen: sternum VIII covered with pale cream-coloured scales.

Pupa (Fig. 1; Table 1). Position and development of setae as figured; range and modal number of branches in Table 1. Measurements and counts from 20 to 22 specimens. Integument without distinctive colour pattern, mostly light brown to yellowish; sterna II–V with narrow dark band near anterior margin. Cephalothorax: setae 1–3-CT about equal in length, 5-CT with 3–8 branches, 6,10,11-CT with 2–4 branches, 7-CT usually with 4 branches (3–6), 8-CT single to triple, 9,12-CT usually triple (2–5). Seta 9-II,III lightly pigmented, 9-IV–VIII pigmented light brown, slightly darker at base. Abdomen: seta 6-I usually triple (2–4), 7-I with 3–9 branches; 1-II dendritic, with > 30 fine branches arising from strong basal stem, 6-II double or triple, 7-II with 4–11 branches, 8-II always absent, 9-II very short, stout, arising cephalad from posterior margin of segment, length 0.01–0.014 mm (mean 0.011 mm), 10,11-II rarely present; 1-III with 5–10 branches, 5-III with 6–11 branches, 6-III usually double (2,3), 9-III short, stout, length 0.01–0.03 mm (mean 0.02 mm); 1-IV with 3–7 branches, 5-IV with 4–8 branches, 6-IV single to triple, 9-IV short, length 0.02–0.05 mm (mean 0.04 mm), ratios of length of 9-IV/9-III 1.25–4.0 (mean 2.36) and 9-IV/9-V 0.19–0.52 (mean 0.36); 1-V usually triple (2–4), 5-V with 3–5 branches, 6-V usually single (1,2), 9-V long, simple, length 0.08–0.12 mm (mean 0.10 mm); 1-VI usually triple (2,3), 5-VI with 3–7 branches, 6-VI single to triple, 9-VI long, frequently simple, occasionally with minute denticles, length 0.09–0.13 mm (mean 0.11 mm); 1-VII double or triple, 5-VII with 4–7 branches, 6-VII usually double (1,2), 9-VII long, frequently simple, sometimes with minute denticles, length 0.09–0.12 mm (mean 0.11 mm); 9-VIII with 12–20 branches. Paddle: lightly tanned, buttress slightly darker, midrib faint, outer basolateral serrations prominent, filamentous spicules on outer apical margin and most of inner margin prominent; seta 1-Pa strong, darkly pigmented, 2-Pa with 1–3 branches; toothed margin index 0.86–0.94 (mean 0.91), paddle teeth well developed, teeth broad at base tapering to blunt apex and widely spaced from each other.

Larva (Fig. 1, Table 2). Position and development of setae as figured; range and modal number of branches in Table 2. Measurements and counts from 10 to 24 specimens. Head: integument light brown to yellowish either with or without pattern of darkened spots, if dark spots present, as follows: 1 dark spot placed posteriorly on dorsal apotome; 2 lateral spots on dorsal apotome posterior to seta 8-C, posteriorly on frontal ecdysial line; 1 dark spot ventral on lateralia encircling seta 12-C; 2 dark spots on labiogula posterior to hypostomal suture; head length 0.60–0.70 mm (mean 0.67 mm), width 0.63–0.74 mm (mean 0.68 mm), antenna length 0.25–0.30 mm (mean 0.28 mm), ratio of distance between base and seta 1-A to antenna length 0.31–0.42 (mean 0.38), seta 2-C long, usually single (1,2), with minute sparse spicules on distal 0.5, 3-C single, length 0.07–0.09 mm (mean 0.08 mm), distance between bases of 2-C and 3-C 0.03–0.04 mm (mean 0.04 mm), 4-C posterolaterad of 2-C, single, length 0.04–0.07 mm (mean 0.06 mm), never reaching base of 2-C, distance between insertions of 2-C and




Fig. 1. *Anopheles latens* sp.n. Pupa (male): CT, cephalothorax; GL, genital lobe; Pa, paddle; I-IX, abdominal segments of pupa (dorsal on left, ventral on right), numbers on the left side denote seta 9 of designated segments. Larva: C, head; P, prothorax; M, mesothorax; T, metathorax; abdominal segments I-VI left side dorsal, right side ventral; VII-X lateral (left side) view. Millimetre scale bars.

Table 1. Number of branches for pupal setae of *Anopheles latens*: range (mode). n.c. = not counted.

Seta no.	Cephalothorax CT	Abdominal segments									Paddle Pa
		I	II	III	IV	V	VI	VII	VIII	IX	
0	–	–	I	I	I	I	I	I	I	–	–
1	2–5 (3)	n.c.	13–26	5–10 (6)	3–7 (5)	2–4 (3)	2,3 (3)	2,3 (2)	–	I	I
2	2,3 (2)	4–9 (6)	4–8 (6)	4–7 (6)	2–6 (4)	2–5 (4)	2–5 (4)	3–6 (4)	–	–	1–3 (2)
3	3–6 (4)	1,2 (1)	2–4 (3)	2–5 (3)	5–11 (6)	1–5 (4)	1–3 (3)	2–5 (4)	–	–	–
4	2,3 (3)	4–8 (6)	4–8 (7)	3–7 (6)	2–5 (4)	3–8 (5)	2–4 (3)	1–3 (3)	1–3 (2)	–	–
5	3–8 (6)	3,4 (3)	3–6 (4)	6–11 (8)	4–8 (5)	3–5 (5)	3–7 (5)	4–7 (6)	–	–	–
6	2–4 (3)	2–4 (3)	2,3 (2)	2–4 (2)	1–3 (2)	1,2 (1)	1–3 (2)	1,2 (2)	–	–	–
7	3–6 (4)	3–9 (5)	4–11 (5)	2–6 (3)	2–5 (4)	2–6 (4)	1,2 (1)	1,2 (1)	–	–	–
8	1–3 (1)	–	–	2–4 (3)	2,3 (2)	2–4 (2)	2,3 (3)	2–4 (3)	–	–	–
9	2–5 (3)	I	I	I	I	I	I	I	12–20 (14)	–	–
10	2–4 (2)	–	0–3	2–5 (3)	1,2 (2)	2,3 (2)	1–3 (2)	2–4 (3)	–	–	–
11	2–4 (2)	–	0–2	I	I	1,2 (1)	1,2 (1)	1–4 (2)	–	–	–
12	2–5 (3)	–	–	–	–	–	–	–	–	–	–
13	–	–	–	–	–	–	–	–	–	–	–
14	–	–	–	–	I	I	I	I	I	–	–

Table 2. Number of branches for setae of fourth-instar larva of *An. latens*: range (mode).

Seta no.	Head C	Thorax			Abdominal segments									
		P	M	T	I	II	III	IV	V	VI	VII	VIII	X	
0	1	1	—	—	—	1	1	1	1	1	1	1	—	
1	1	10–17 (13)	22–32 (28)	2–3 (3)	3–7 (6)	8–14 (10)	11–20 (18)	13–21 (18)	15–21 (19)	14–19 (16)	11–17 (12)	1	1	
2	1–2 (1)	9–13 (12)	2–5 (3)	1	2–3 (2)	5–9 (7)	5–9 (6)	3–5 (3)	3–5 (4)	4–7 (5)	4–8 (6)	6–10 (7)	17–21 (17)	
3	1	1	1	8–11 (9)	1–2 (1)	1	1	1–3 (2)	1	1	3–4 (3)	6–10 (8)	7–11 (8)	
4	1	13–20 (16)	2–4 (3)	2–4 (3)	3–6 (5)	3–9 (6)	3–5 (3)	3–5 (3)	3–4 (3)	1	1–2 (1)	1	9	
5	7–11 (9)	25–44 (31)	1	21–38 (34)	3–4 (3)	3–6 (5)	1–5 (3)	3–4 (3)	3–4 (3)	3–5 (4)	4–6 (5)	3–7 (3)	—	
6	8–14 (12)	1	3–4 (3)	2–4 (3)	14–21 (17)	18–29 (21)	10–19 (14)	1–3 (2)	1–3 (2)	2–4 (3)	3–5 (3)	1–S	3–5 (5)	
7	9–16 (12)	15–26 (21)	1–4 (2)	20–32 (30)	11–19 (13)	17–24 (20)	3–6 (4)	4–6 (5)	2–5 (4)	2–5 (3)	1–6 (4)	2–S	4–7 (6)	
8	1–3 (2)	19–29 (27)	12–21 (16)	16–26 (21)	—	1–3 (2)	1–3 (2)	2–3 (3)	2–3 (2)	2–3 (3)	3–5 (4)	6–S	1–3 (2)	
9	2–4 (4)	1–2 (2)	1–2 (1)	1–2 (1)	3–5 (4)	6–8 (7)	6–11 (9)	5–12 (8)	5–8 (6)	5–7 (6)	4–6 (5)	7–S	1	
10	1–2 (2)	1	1	1	1	2–4 (3)	1	1	1	2–4 (3)	4–7 (6)	8–S	2–4 (3)	
11	18–30 (21)	1	1	1	3–4 (3)	1	2–4 (3)	2–4 (3)	2–4 (3)	1–3 (3)	1–3 (2)	9–S	3–5 (4)	
12	1–2 (2)	1	1	2–3 (2)	2–4 (3)	1	2–4 (3)	2–5 (3)	1–4 (3)	1	1–2 (1)	—	—	
13	3–6 (6)	3–6 (5)	3–7 (5)	2–4 (3)	2–6 (4)	3–5 (3)	4–7 (5)	3–5 (3)	3–5 (4)	5–8 (7)	2–4 (3)	—	—	
14	3–5 (5)	5–8 (6)	5–11 (8)	—	—	—	—	1	1	1	1	1	—	
15	3–5 (5)	—	—	—	—	—	—	—	—	—	—	—	—	

4-C 0.06–0.10 mm (mean 0.08 mm), ratio of length of 4-C to distance between insertions of 2-C and 4-C 0.51–0.88 (mean 0.74), distance between bases of 3-C and 4-C 0.06–0.09 mm (mean 0.07 mm), 5-C longer than antenna, extending beyond anterior margin of head, with 7–11 branches, 6-C with 8–14 branches, 7-C with 9–16 branches. Thorax: tubercles of all large setae light brown; seta 1-P with 10–17 branches, stem stout, not noticeable expanded, flattened basally, arising from large tubercle, tubercle frequently separated from that of 2,3-P or rarely both tubercles joined at base, each tubercle with small, inconspicuous apical lip arising from posterodorsal side and projecting forward over bases of 1,2-P, 14-P with 5–8 branches; 4-M with 2–4 branches, 6-M usually triple (3,4), 14-M with 5–11 branches; 3-T weakly developed, palmate, with 8–11 narrow lanceolate nearly transparent leaflets. Abdomen: seta 1-I small, somewhat palmate, with 3–7 narrow lanceolate nearly transparent leaflets, 2-I double or triple, 3-I usually single (1,2), 9-I with 3–5 branches; 1-II small, less pigmented than 1-I, somewhat palmate, with 8–14 narrow lanceolate nearly transparent leaflets, basal stem narrow, poorly pigmented; 2-IV with 3–5 branches, 3-IV single to triple, 13-IV long, with 3–5 branches, ratio of length of 13-IV to 10-IV 0.46–0.64 (mean 0.55); 2-V with 3–5 branches; 1-VII smaller and less pigmented than 1-VI, with 11–17 moderately broad lanceolate leaflets with minute apicolateral serrations and apical filament; 1-X long, single, inserted on saddle; pecten with 4–6 long teeth alternating with 6–10 short teeth, with 11–15 total teeth.

Material examined. HOLOTYPE: adult male with associated larval and pupal exuviae on microscope slide, acc. no. 08161–10, collected by AFRIMS team, 25 May 1980, deposited in the NMNH. Type locality: THAILAND, Phangnga, Ban Bang Kaeo (8°35'N 98°32'E). PARATYPES: INDONESIA, Kalimantan, Salaman (3°49'S 115°5'E), 19–24 September 1986, acc. no. IDK 43, 1F, 1M, 2Le, 2Pe. MALAYSIA (EAST), Sabah, Beaufort (5°20'N 115°45'E), 4 April 1970, acc. no. S-368, 1M, 1Le, 1Pe. THAILAND, Chumphon, Pathiu, Ban Chong Rakam, Ban Chong Mut, Mu 3 (10°44'N 99°18'E), 18 September 1978, acc. no. 08007(3), progeny, 1F, 1M, 3Le, 3Pe. Nakhon Si Thammarat, Tung Song, Nam Tok (8°10'N 99°30'E), 1985, acc. no. TS39 (F-2), progeny, 1F, 3M, 5Le, 5Pe, 3I; Tung Yai, Ban Tham Phae Dan (8°22'N 99°20'E), 3–8 December 1985, acc. no. TY001, progeny, 2F, 2M, 4Le, 5Pe. Phangnga, Ban Bang Kaeo (8°35'N 98°32'E), 25 May 1980, acc. no. 08161, 6F, 8M, 11Le, 11Pe; 25 May 1980, acc. no. 08162, 3F, 2M, 5Le, 5Pe. Songkhala: Sadao, Padang Besa, Khao Rup Chang (6°40'N 100°19'E), 12 December 1986, acc. no. PB53, progeny, 4F, 1M, 4Le, 5Pe. Paratypes deposited in the NMNH. Total material of *An. latens* examined: 263 specimens comprising 73F, 43M, 55Le, 69Pe, 23 L derived from 52 separate collections from natural habitats (28 adults, 24 immatures) and 10 progeny broods.

Distribution. *Anopheles latens* is known from Indonesia (Kalimantan), east Malaysia (Sabah, Sarawak), west Malaysia (Malay States, Kelantan, Negeri Sembilan, Pahang, Perak,

Perlis and Selangor) and Thailand (Chumphon, Nakhon Si Thammarat, Narathiwat, Phangnga, Satun and Songkhala).

Bionomics. The importance of *An. latens* as vector of human malaria parasites, and bancroftian filariasis parasites is not well established because this species has been largely misidentified as *An. leucosphyrus*. Probably most of the literature records regarding *An. leucosphyrus* from Indonesia, east and west Malaysia and Thailand are *An. latens*. Based on the published records for *An. leucosphyrus* from the localities mentioned before, *An. latens* seems to be an important vector of human malaria parasites and also involved in the transmission of bancroftian filariasis in Sarawak (Zulueta & LaChance, 1956; Zulueta, 1956, 1957). In a mountainous area in the tropical rain forest of Sarawak, *An. latens* was found to be more abundant in farm huts than in village settlements. The malaria transmission was more intense in former than in latter, but in both ecotypes, females were attracted more to humans than to CDC light traps. The inoculation rate of *An. latens* in farm huts was 0.023 (Seng et al., 1999). Also, *An. latens* was incriminated as a vector of both human malaria and Bancroftian filarial parasites in villages and in forested areas situated 0.5 km from the villages. Females activity peaked around midnight in forested areas and soon after dusk in village settlements (Chang et al., 1995). These differences in the peak activity may be related to changes in the microenvironmental conditions in both ecotypes, thus the humidity is lower indoors than outdoors during the night and the opposite in the dusk. Human bait collections at a remote village located in a heavily forested area in south Kalimantan, Indonesia, showed that *An. latens* and *An. balabacensis* were predominant outside houses in villages, whereas the number was lower in nearby forest. The sporozoite infection rate was 1.0% for *An. latens*, slightly less than 1.3% for sympatric *An. balabacensis* (Harbach et al., 1987).

Like other members of the *Leucosphyrus* Complex, *An. latens* breeds in forests where larval habitats are mostly in shaded temporary pools and natural containers on the ground. In west Malaysia, immatures were found in a muddy pool in a cart track running through dense jungle and in the longitudinal half of a bamboo stem on the ground (Leicester, 1903), and in jungle streams in Selangor, Jugra (Leicester, 1908a).

Collections in Thailand for this study included immatures of *An. latens* from ground pools along stream margins, flood pools, seepage pools, sandy pools in stream banks, small shallow running streams and elephant footprints. The water was either fresh or stagnant, turbid or clear, sometimes with some degree of pollution and decaying leaves. The habitats were partially or heavily shaded in primary and secondary rain forests of mountains or valleys at elevations ranging from 76 to 520 m above sea level. Immatures were found in association with those of many other mosquito species: *Aedes orbitae* Edwards, *Ae. vexans* (Meigen), *Anopheles baimaii* sp.n., *An. bengalensis* Puri, *An. introlatus* Colless, *An. macarthurii* Colless, *An. maculatus* Theobald, *An. pujutensis* Colless, *An. nemophilous* Peyton & Ramalingam, *Culex* spp. including *Cx. fraudatrix* (Theobald),

Cx. nigropunctatus Edwards, *Cx. mimulus* Edwards and *Cx. minor* (Leicester), *Verrallina atria* (Barraud) and *Uranotaenia* spp. including *Ur. bicolor* Leicester and *Ur. trilineata* Leicester.

Adults of *An. latens* were collected on human bait in secondary evergreen forest, and secondary rain forest with fruit and/or rubber plantations and/or orchards near villages and/or mixed fruit plantations. Collections were made at the edge of forest or near human habitations in villages, valleys or mountainous or hilly areas.

Immatures were found in swamps in east Malaysia. The water was static, fresh and clear. Larval habitats were situated in partial shade in secondary rain forest in hilly areas. In west Malaysia, larval habitats were rock pools situated on the sides of running streams, flood pools, ground pools and large swamps. The water was fresh, stagnant, and clear, sometimes with decaying leaves. The habitats were found in partial shade in secondary and primary rain forests situated in valley, mountainous or hilly areas. Immatures were found in association with *Ae. caecus* (Theobald), *Ae. varietus* Doleschall, *An. hodgkini* Reid, *An. scanloni* sp.n., *Cx. foliatus* Brug., *Cx. mammilifer* (Leicester), *Cx. mimulus*, *Ur. bicolor* and *Verrallina gibbosa* (Delfinado). Adults were collected on human bait at ground level as well as in forest canopy.

Systematics. The *Leucosphyrus* Complex includes four species: *An. balabacensis*, *An. introlatus*, *An. latens* (as *An. leucosphyrus* A) and *An. leucosphyrus* s.s. (Peyton, 1990). Based on hybridization and cytogenetic evidence, Kanda *et al.* (1981) suggested that specimens from populations of Sabah, and Kuching (SWK) and Niabet (SWN) in Sarawak, Malaysia, belonged to a separate species they termed '*An. leucosphyrus sensu stricto*'. Later, Kanda *et al.* (1983) using chromosomal polymorphisms confirmed that both SWK and SWN populations belonged to their '*An. leucosphyrus* s.s.' Using seven protein loci, Takai (1986) was able to distinguish five taxa within the *An. leucosphyrus* group and confirmed the hypothesis of Kanda *et al.* (1981, 1983) that specimens of '*An. leucosphyrus*' from SWK belonged to a distinct species. Finally, Baimai *et al.* (1988b) found cytogenetic evidence that *An. leucosphyrus* s.l. included two allopatric species, one inhabiting Borneo, west Malaysia and southern Thailand (designated as *An. leucosphyrus* A), the other confined to Sumatra (*An. leucosphyrus* s.s.). Based on Baimai *et al.*'s findings, we conclude that specimens termed '*An. leucosphyrus* s.s.' by Kanda *et al.* (1981, 1983) and Takai (1986) were *An. latens*.

Anopheles latens is morphologically more similar in all life stages to *An. leucosphyrus* than to any other species of the *Leucosphyrus* Complex. The adult males and females of *An. latens* are indistinguishable from those of *An. leucosphyrus*; however, they can be separated from *An. introlatus* and *An. balabacensis* by having the PSD spot of vein R extending basally to or beyond the level of the HD spot on the costa, from level 4 to level 6, and the apical pale band of palpmere 5 distinctly cream-coloured or yellowish contrasting with the silvery white bands of palpmeres 2 and 3. In *An. introlatus* and *An. balabacensis*, the PSD spot of vein R usually does not extend basally beyond the level of the PSD spot on the costa (level 1) and rarely extends into

the distal 0.5 of the HD spot of the costa (level 4), but never reaches or extends beyond level 5; the apical pale band of palpmere 5 is white or whitish, not strongly contrasting with the silvery white bands of palpmeres 2 and 3. The pupal stage of *An. latens* cannot be distinguished from those of *An. balabacensis* and *An. introlatus* but some individuals can be distinguished from that of *An. leucosphyrus* by having seta 7-II with 4–11 branches, seta 2-VII with 3–6 branches and 2-V with 2–5 branches, whereas in *An. leucosphyrus* seta 7-II 2–6 branches, 2-VII 2,3 branches and 2-V always triple. Although useful to separate some individuals of these two species, the pupal characters overlap in *An. leucosphyrus* and *An. latens*, thus they should be used with caution for species separation. The fourth-instar larva of *An. leucosphyrus* and *An. latens* can be distinguished from those of *An. balabacensis* and *An. introlatus* by having the tubercles of seta 1-P and 2-P without a posterodorsal tooth, or at most with a weak and inconspicuous lip projecting forward over the base of each seta, and with the tubercles of both setae usually well separated, occasionally joined basally. In *An. balabacensis* and *An. introlatus*, the basal tubercle of setae 1-P and 2-P are joined basally, and each tubercle has a prominent tooth or spine arising from the posterodorsal margin that projects over the base of the seta. The fourth-instar larvae of *An. latens* and *An. leucosphyrus* are indistinguishable. Since these two species are allopatric, their distributions should be considered when making identifications.

Dirus complex

Anopheles (Cellia) cracens Sallum & Peyton, sp.n.

Etymology: the name *cracens* is Latin for neat, graceful.

Anopheles balabacensis Perlis/Thai Form of Reid (1968) (taxonomy).

Anopheles balabacensis Perlis Form of Baimai *et al.*, 1981: 81 (cytogenetics, chromosome differentiation, evolution); Sucharit & Choochote, 1983: 90 (morphometry of male genitalia, frequency of gonostylus movement).

Anopheles balabacensis perlisiensis of Yong *et al.*, 1983: 611 (electromorphs, species differentiation); Ward, 1984: 260 (*nomen nudum*).

Anopheles dirus B of Baimai *et al.*, 1984: 536 (metaphase chromosomes); Wibowo *et al.*, 1984: 425 (H-banding pattern of sex chromosomes); Peyton, 1990: 197 (taxonomy); Klein *et al.*, 1991: 481 (susceptibility to *P. cynomolgi* B strain); Prapanthadara *et al.*, 2000: 395 (glutathione S-transferase).

Anopheles dirus species B of Hii, 1984a: 104 (DDT/fenitrothion tolerance); Hii, 1984b: 192 (crossbreeding); Hii, 1985: 185 (polytene chromosomes, crossbreeding); Hii, 1986: 125 (morphometry L, P); Baimai & Traipakvasin, 1987: 401 (heterochromatin sex chromosome); Baimai, 1988a: 13 (heterochromatin sex chromosomes, distribution, evolution); Baimai, 1988b: 667 (cytogenetics, distribution,

bionomics); Baimai *et al.*, 1988d: 372 (polytene chromosomes); Baimai *et al.*, 1988a: 333 (crossbreeding, heterochromatin); Baimai *et al.*, 1988c: 151 (bionomics, distribution); Yasothornsrikul *et al.*, 1988: 703 (RFLP); Damrongphol & Baimai, 1989: 563 (egg morphology); Green *et al.*, 1992: 29 (enzyme electromorphs); Audtho *et al.*, 1995: 107 (DNA hybridization); Poopittayasatporn & Baimai, 1995: 426 (polytene chromosome, phylogeny); Sithiprasasna *et al.*, 1996: 483 (methoprene resistance); Walton *et al.*, 1999: 24 (ASA PCR species identification); Huang *et al.*, 2001: 615 (PCR species identification); Manguin *et al.*, 2002: 46 (SCAR multiplex PCR, species identification).

Anopheles leucosphyrus Perlis (IMR) of Kanda *et al.*, 1981: 321 (hybridization, phylogeny).

Female. Head: proboscis length 1.97–2.25 mm (mean 2.11 mm), ratio of length to forefemur 1.09–1.17 (mean 1.13), maxillary palpus length 1.82–2.10 mm (mean 1.96 mm), ratio of length to proboscis length 0.89–0.97 (mean 0.93), ratio of length to forefemur length 1.01–1.12 (mean 1.05), ratio of length of palpomeres 3/4 1.67–1.92 (mean 1.78), 3/5 2.18–2.78 (mean 2.45), 4/5 1.20–1.56 (mean 1.37), 4–5/3 0.88–1.05 (mean 0.97), palpomeres 2–4 with narrow apical silvery white bands, pale scales of palpomere 5 white to pale cream-coloured, length of apical pale band of palpomere 5 1.43–3.67 (mean 2.12) length of basal dark band of palpomere 5, ratio of length of apical pale band of palpomere 4 to length of basal dark band of palpomere 5 0.50–1.25 (mean 0.88). Thorax: pleural setae as follows: 1–3 upper proepisternal, 1–3 prespiracular, 3–5 prealar, 3–6 upper mesokatepisternal, 2,3 lower mesokatepisternal, 3–5 upper mesepimeral. Wing: length 3.21–3.74 mm (mean 3.45 mm), pale scales on all veins light cream-coloured, spots on costa and subcosta slightly more obviously cream-coloured, not strongly contrasting with other spots; PHP spot of costa always present, prominent or reduced, HP spot always present, small, PSP and SP spots always present, prominent, ASP spot always absent, PP spot 0.75–1.44 (mean 1.14) length of SCP spot, AD spot 1.56–3.00 (mean 2.13) length of PP spot, PSD spot of vein R extending basally from level 1 to level 5 on one or both wings, PSD spot of vein R with 1–3 small pale interruptions, SD spot of vein R with 1–5 pale interruptions, number of pale interruptions on PSD-PD spots of vein R 4–9 on each wing, ratio of length of cell R_2 to vein R_{2+3} 1.33–1.95 (mean 1.66), ratio of length of cell R_2 to cell M_{1+2} 1.23–1.38 (mean 1.30). Legs: foretarsomeres 2 and 3 with prominent basal and apical pale spots, dark area of foretarsomere 2 with or without 1–3 pale spots on dorsal surface mostly pale-scaled on dorsal surface with small dark median band, mostly dark-scaled on ventral surface, foretarsomere 3 with apical and basal pale bands, bands sometimes poorly developed, foretarsomere 4 noticeably pale-scaled at base, apical pale band usually prominent, rarely inconspicuous, foretarsomere 5 mostly dark-scaled with pale scales at apex, foretarsomeres 1–5 mostly dark-scaled on ventral surface, basal and apical pale bands of ventral surface absent or present but less evident than those on dorsal surface; midtarsomere

2 usually without basal pale band, rarely with inconspicuous basal pale band, median dark region usually with 1–3 pale spots, rarely entirely dark-scaled, basal and apical pale bands complete or incomplete, midtarsomere 3 with apical pale band more distinct on dorsal surface, midtarsomeres 4 and 5 with complete or incomplete apical pale bands, sometimes mostly pale-scaled on ventral surface; hindtarsomere 2 with apical pale band, basal pale band absent or inconspicuous, median dark part entirely dark-scaled or with 1–7 small pale spots, hindtarsomere 3 with apical pale band, hindtarsomere 4 with apical and complete or incomplete basal pale bands, hindtarsomere 5 usually without basal pale scales, sometimes with inconspicuous basal pale band, apical pale band always present. Abdomen: tergum VI usually with 2–5 pale cream-coloured to tan scales on posterolateral margin, tergum VII variable, with few or occasionally several posterolateral scales, scales vary from pale cream-coloured to tan to dark, tergum VIII with a large posteromedial patch of pale cream-coloured to golden scales; sternum VI with small patch of dark scales posteromedially, sternum VII with posteromedial patch of dark scales, rarely with 2,3 lateral pale scales, sternum VIII with small anterolateral patch of whitish scales.

Male. Essentially as in female except for sexual characters. Wing: generally paler with reduced scaling, pale spots usually longer than in female. Palpomere 2 with dorsal patch of pale scales at middle, extending onto lateral surface, apex of palpomere 2 bare; palpomere 3 with long median dorsal patch of pale scales at middle, extending onto lateral surface, apex of palpomere 3 with broad band of pale scales covering dorsal, lateral and ventral surfaces except for narrow ventrolateral line of dark scales; palpomeres 4 and 5 mostly pale-scaled with narrow basal band of dark scales, basal dark band of palpomere 5 with short extension on ventral surface not reaching apex. Abdomen: sternum VIII covered with pale cream-coloured scales.

Pupa (Fig. 2, Table 3). Position and development of setae as figured; range and modal number of branches in Table 3. Measurements and counts from 20 to 75 specimens. In general similar to *An. latens* except for following characters. Sterna II–VII with narrow dark band near anterior margin. Cephalothorax: 5-CT with 5–7 branches, 8,10-CT normally double (1–3), 9,11,12-CT usually triple (9, 2–4; 11,12, 2–5). Seta 9-II–VII very lightly pigmented, 9-VIII light brown, slightly darker at base. Abdomen: seta 6-I usually double (1–3), 7-I with 3–7 branches; 6-II single to triple, 7-II with 3–7 branches, 8,10,11-II absent, 9-II length 0.01–0.02 mm (mean 0.01 mm); 1-III with 8–17 branches, 5-III with 7–15 branches, 6-III usually double, 9-III length 0.01–0.03 mm (mean 0.02 mm); 1-IV with 4–13 branches, 5-IV with 6–14 branches, 6-IV always single, 9-IV long, usually with small spines, length 0.04–0.11 mm (mean 0.08 mm), ratios of length of 9-IV/9-III 2.00–6.86 (mean 4.35) and 9-IV/9-V 0.41–0.96 (mean 0.78); 1-V with 2–6 branches, 5-V with 5–9 branches, 9-V usually with minute denticles, length 0.09–0.12 mm (mean 0.11 mm); 1-VI with 2–4 branches, 5-VI with 4–8 branches, 6-VI single, 9-VI usually with minute denticles, length 0.10–0.13 mm (mean 0.11 mm); 1-VII single




Fig. 2. *Anopheles cracens* sp.n. Pupa (male): CT, cephalothorax; GL, genital lobe; Pa, paddle; I-IX, abdominal segments of pupa (dorsal on left, ventral on right), numbers on the left side denote seta 9 of designated segments. Larva: C, head; P, prothorax; M, mesothorax; T, metathorax; abdominal segments I-VI left side dorsal, right side ventral; VII-X lateral (left side) view. Circle shows basal details of setae 1-3-P. Millimetre scale bars.

Table 3. Number of branches for pupal setae of *Anopheles cracens*: range (mode). n.c. = not counted.

Seta no.	Cephalothorax CT	Abdominal segments									Paddle P
		I	II	III	IV	V	VI	VII	VIII	IX	
0	-	-	1	1	1	1	1	1	1	-	-
1	2-4 (2)	n.c.	>30	8-17 (11)	4-13 (7)	2-6 (3)	2-4 (2)	1-3 (2)	-	1	1
2	2-3 (2)	3-9 (6)	4-8 (5)	4-8 (6)	3-4 (3)	2-4 (3)	3-4 (3)	3-4 (3)	-	-	1-2 (1)
3	3-4 (3)	1	2-4 (3)	3-5 (3)	5-9 (7)	3-6 (3)	2-3 (2)	3-4 (3)	-	-	-
4	3-4 (4)	4-8 (6)	5-9 (5)	3-5 (5)	3-5 (4)	3-7 (5)	1-3 (3)	1-2 (2)	2-3 (2)	-	-
5	5-7 (5)	3-5 (3)	3-6 (5)	7-15 (10)	6-14 (8)	5-9 (7)	4-8 (6)	4-8 (5)	-	-	-
6	2-4 (3)	1-3 (2)	1-3 (2)	1-3 (2)	1	1-2 (1)	1	1-2 (2)	-	-	-
7	3-5 (4)	3-7 (5)	3-7 (4)	2-5 (3)	2-6 (3)	2-5 (3)	1-2 (1)	1-2 (2)	-	-	-
8	1-3 (2)	-	-	2-3 (2)	1-4 (3)	2-3 (3)	1-3 (2)	2-4 (3)	-	-	-
9	2-4 (3)	1	1	1	1	1	1	1	12-18 (16)	-	-
10	1-3 (2)	-	-	2-5 (3)	1-3 (2)	1-3 (2)	1-4 (2)	2-4 (2)	-	-	-
11	2-5 (3)	-	-	1-2 (1)	1	1	1	1-2 (2)	-	-	-
12	2-5 (3)	-	-	-	-	-	-	-	-	-	-
13	-	-	-	-	-	-	-	-	-	-	-
14	-	-	-	-	1	1	1	1	1	-	-

Table 4. Number of branches for setae of fourth-instar larva of *An. cracens*: range (mode).

Seta no.	Head C	Thorax			Abdominal segments								
		P	M	T	I	II	III	IV	V	VI	VII	VIII	X
0	1	1	-	-	-	1	1	1	1	1	1	1	-
1	1	7-20 (15)	25-29 (27)	1	4-10 (9)	5-17 (5)	15-21 (21)	13-20 (19)	11-21 (20)	14-18 (17)	7-13 (11)	1	1
2	1	9-12 (11)	1-3 (1)	1	1-3 (2)	6-8 (8)	6-8 (6)	3-5 (4)	3-5 (4)	4-5 (5)	4-7 (6)	4-7 (5)	15-23 (22)
3	1	1	1	3-6 (4)	1-3 (2)	1	1	2-3 (2)	1	1	2-3 (3)	5-7 (6)	9-14 (9)
4	1	13-19 (15)	2-3 (3)	2-5 (4)	3-7 (4)	4-7 (5)	3-4 (4)	2-4 (3)	3-5 (3)	1	1	1	9
5	7-12 (11)	32-38 (38)	1-2 (1)	24-40 (29)	3-6 (3)	3-7 (5)	3-5 (4)	3-5 (3)	3-5 (3)	4-5 (5)	3-5 (3)	3-4 (3)	-
6	9-16 (13)	1	2-5 (4)	2-3 (3)	15-23 (20)	15-29 (23)	12-22 (14)	1-3 (2)	1-3 (2)	2-3 (3)	2-4 (3)	1-S	4-6 (5)
7	11-16 (11)	17-27 (24)	2-3 (3)	24-35 (29)	15-23 (16)	20-26 (20)	3-5 (4)	4-6 (5)	4-6 (4)	2-5 (3)	3	2-S	4-7 (4)
8	1-2 (1)	21-32 (30)	18-29 (25)	23-35 (27)	-	1-2 (1)	1-2 (1)	2-3 (2)	2-3 (2)	2-3 (2)	3-8 (3)	6-S	2-3 (2)
9	1-5 (2)	1	1	1	3-5 (4)	5-9 (8)	6-9 (9)	7-12 (9)	7-9 (8)	6-8 (7)	4-5 (4)	7-S	1-2 (2)
10	1-2 (1)	1	1	1	1	1-3 (3)	1	1	1	1-4 (3)	4-7 (5)	8-S	2
11	24-40 (31)	1	1	1	3-4 (3)	1	2-4 (3)	2-3 (3)	2-3 (2)	2-3 (2)	2	9-S	2-4 (3)
12	1-2 (2)	1	1	1-3 (2)	2-5 (3)	1-3 (3)	2-3 (2)	2-3 (2)	2-3 (3)	1	1	-	-
13	4-7 (6)	4-6 (4)	2-5 (5)	2-3 (3)	2-5 (4)	3-6 (4)	3-7 (5)	3-4 (3)	3-4 (3)	3-8 (6)	2-3 (2)	-	-
14	4	4-8 (6)	4-8 (7)	-	-	-	-	1	1	1	1	1	-
15	4-7 (7)	-	-	-	-	-	-	-	-	-	-	-	-

to triple, 5-VII with 4–8 branches, 9-VII usually with minute denticles, length 0.09–0.13 mm (mean 0.11 mm); 9-VIII with 12–18 branches. Paddle: seta 2-Pa single or double; toothed margin index 0.80–0.92 (mean 0.87), teeth well developed, strong, widely separated.

Larva (Fig. 2, Table 4). Position and development of setae as figured; range and modal number of branches in Table 4. Measurements and counts from eight to 29 specimens. In general as described for *An. latens* except for following characters. Head: integument very lightly pigmented, without pattern of dark spots; length 0.59–0.77 mm (mean 0.70 mm), width 0.66–0.82 mm (mean 0.75 mm); antenna length 0.26–0.32 mm (mean 0.29 mm), ratio of distance between base and seta 1-A to antenna length 0.30–0.53 (mean 0.40); seta 3-C length 0.07–0.10 mm (mean 0.08 mm); distance between bases of 2-C and 3-C 0.03–0.05 mm (mean 0.04 mm); 4-C length 0.05–0.10 mm (mean 0.07 mm), either extending to or not reaching base of 2-C, distance between insertions of 2-C and 4-C 0.06–0.10 mm (mean 0.08 mm), ratio of length of 4-C to distance between insertions of 2-C and 4-C 0.73–1.09 (mean 0.92); distance between base of 3-C and 4-C 0.06–0.09 mm (mean 0.07 mm); 5-C longer than antenna, not reaching anterior margin of head, with 7–12 branches, 6-C with 9–16 branches, 7-C with 11–16 branches. Thorax: tubercles of all large setae very lightly pigmented, light tan to pale straw-yellow; seta 1-P with 7–20 branches, tubercle of 1-P joined to tubercle of 2,3-P, each tubercle with a strong apical tooth arising from posterodorsal side and projecting forward over bases of 1,2-P, 14-P with 4–8 branches; 4-M double or triple, 6-M 2–5-branched, 14-M with 4–8 branches; 3-T moderately developed, palmate, with 3–6 leaflets. Abdomen: seta 1-I with 4–10 leaflets, 2,3-I single to triple; seta 1-II more developed than 1-I, palmate, with 5–17 leaflets; 3-IV double or triple, 13-IV with 3,4 branches, ratio of length of 13-IV to 10-IV 0.54–0.79 (mean 0.65); 1-VII with 7–13 leaflets; 1-X long, single, inserted in marginal notch or at edge of saddle; 11–16 pecten teeth, 3–5 long teeth alternating with 7–12 short teeth.

Material examined. HOLOTYPE: adult female with associated larval and pupal exuviae on microscope slide, acc. no. MH0023(1)–4, collected by V. Baimai and R. G. Andre, 28 April 1982, from progeny brood, deposited in the NMNH. Type locality: MALAYSIA (WEST), Terengganu, Kampong Tapah (5°6'N 102°55'E). PARATYPES: MALAYSIA (WEST): Terengganu: Kampong Dura (5°4'N 102°56'E), 26 April 1982, collection (accession) number MH0008 (1), progeny, 3F, 5M, 8Le, 8Pe; same data, 28 April 1982, progeny, acc. no. MH0016 (1), 6F, 4M, 10Le, 10Pe, 1 L; Kampong Tapah (5°6'N 102°55'E), 28 April 1982, acc. no. 0023 (1), progeny, 5F, 3M, 8Le, 8Pe, 4L. THAILAND: Chumphon: Wang Mai, Ban Jo Po Ro (10°29'N 98°55'E), 20 September 1978, acc. no. 08009, 1F, 2M, 3Le, 3Pe, 1 L. Phangnga: Khao Pak Chaung (8°17'N 98°23'E), 13 October 66, acc. no. 1650, 2M, 2Le, 2Pe. Phattalung: Tanod, Ban Lo Chang Kra (7°10'N 99°30'E), February 1985, acc. no. PT-23 (1), progeny, 4F, 4Le, 4Pe. Paratypes deposited in the NMNH. Total material of *An. cracens* examined: 330 specimens comprising 82F, 39M, 93Le, 95Pe, 21 L, derived from 15

separate collections from natural habitats (12 adult, 3 immature) and 12 progeny broods.

Distribution. *Anopheles cracens* is known from Indonesia (Sumatra Island), west Malaysia (Perlis, Terengganu) and Thailand (Chumphon, Phangnga, Phattalung).

Bionomics. *Anopheles cracens* is an anthropophilic species, showing peak biting activity at 1900–2100 hours and a low level of activity throughout the night outdoors in Phattalung, Thailand (Baimai *et al.*, 1988c). In Sabang Island (Pulau Weh), Aceh, Sumatra, *An. cracens* was captured biting humans at night (J. E. Hudson). In west Malaysia, adult females were also captured in human-bait collections at night in secondary tropical rain forest and in villages in localities situated either in mountainous or hilly areas. Presumably *An. cracens* is an important vector of human malaria, but this requires investigation now it can be identified. A study on the susceptibility to the simian malaria parasite *Plasmodium cynomolgi* B strain in comparison to other South-east Asian *Anopheles* showed that 77% of *An. cracens* were infected (Klein *et al.*, 1991). In another study, *An. cracens* was orally infected with *Brugia pahangi* microfilariae, but the infectivity rate was lower than that of *An. stephensi* Liston (Zahedi & White, 1994). Studies on the susceptibility of a colony of *An. cracens* to DDT, fenitrothion, propoxur and malathion showed a considerable degree of tolerance to the first two chemicals (Hii, 1984a). *Anopheles cracens* was found to be stenogamous in the laboratory (Sucharit & Choochote, 1983) and this facilitated many experimental studies of '*An. dirus* species B' cited above.

In Thailand, immatures were taken from elephant and other animal footprints. The water was fresh, stagnant, temporary, clear or coloured, in full sun or partial shade. The larval habitats were in secondary rain forest situated in both plains and mountainous areas. Immatures of *Ae. caecus*, *Ae. orbitae*, *An. baimaii* sp.n., *An. barbirostris* Van der Wulp and *An. kochi* Dönitz were found in association with *An. cracens*.

Systematics. *Anopheles cracens* was first distinguished from *An. balabacensis* and *An. dirus* by its distinctive mitotic and meiotic karyotypes and the larval salivary gland polytene chromosome banding patterns of individuals from a population designated as *An. balabacensis* Perlis Form (Baimai *et al.*, 1981). Morphometrics of the male genitalia and the frequency of clasper movement also showed differences between *An. cracens* (as *An. balabacensis* Perlis Form) and *An. dirus* (Bangkok colony) (Sucharit & Choochote, 1983). Further examination of the polytene chromosomes and crossbreeding of laboratory colonies by Hii (1985) confirmed the status of *An. dirus* and revealed another species designated as *An. dirus* species B (= *An. cracens*). Additional studies of cytogenetics and hybridization (Baimai *et al.*, 1984, 1988a,d; Hii, 1984b, 1985; Wibowo *et al.*, 1984; Baimai & Traipakvasin, 1987; Baimai, 1988a,b; Poopittayasataporn & Baimai, 1995), plus morphological and morphometric characters of the immature

stages, were taken to confirm the specific status of *An. cracens* within the Dirus Complex (Hii, 1986; Damrongphol & Baimai, 1989). Diagnostic enzyme electrophores, DNA probes and DNA markers have been developed to separate members of the Dirus Complex and to confirm their species status, including *An. cracens* (Yasothornsrikul *et al.*, 1988; Green *et al.*, 1992; Audtho *et al.*, 1995; Walton *et al.*, 1999; Huong *et al.*, 2001; Manguin *et al.*, 2002). Backcrosses of hybrids between *An. dirus* and *An. cracens* showed that autosomes as well as the X chromosome might contribute to sterility (Hii, 1984b, 1985).

Adults of *An. cracens* can be recognized by the following combination of characters: (1) proboscis uniformly dark-scaled, slightly longer than the forefemur, ratio 1.09–1.17 (mean 1.13); (2) PSD spot of vein R with 1–3 small pale interruptions; (3) ASP spot always absent from costa; (4) hindtarsomere 4 always with either a complete or incomplete basal pale band; (5) hindtarsomere 5 usually dark-scaled at base, rarely with an inconspicuous pale band; (6) apical white band of hindtibia without a dark extension into the basal portion; (7) vein 1 A without a noticeably longer pale spot at the level of the PSD spot of the costa; (8) PSD spot of vein R frequently extending basally to level 3 and level 4, less frequently to level 5, and rarely to level 1 and level 2; (9) sternum VI with a small posteromedial patch of dark scales present in both sexes.

Fourth-instar larvae of *An. cracens* can be recognized by the following combination of characters: (1) pigmentation of sclerotized structures very light, inapparent, light tan to pale straw-yellow, the margins of sclerotized tubercles of the larger setae, the anterior and posterior tergal plates, and the saddle not discernable, the head capsule (except for very dark collar), including antenna, uniformly very pale tan to straw-coloured, larger setae light brown, seta I–III–VI slightly darker, small scattered nearly colourless spicules on inner ventral aspect of antenna and on posterolateral surface of saddle; (2) seta 3-C single; (3) seta 5-C longer than antenna; (4) basal sclerotized tubercle of seta 1-P with prominent tooth or spine arising from posterodorsal margin; (4) tubercles of setae 1,2-P broadly joined basally; (5) seta 1-P small and weak, not noticeably expanded basally, with 7–20 branches; (6) seta 4-C short, either not reaching base of 2-C or extending slightly beyond it, ratio of length of 4-C to distance between the insertions of 2-C and 4-C 0.73–1.09; (7) seta 3-C short, extending to or only slightly beyond anterior margin of head; (8) seta 1-II palmate, weakly developed, with 5–17 narrow, lanceolate leaflets; (9) seta 2-IV,V with 3–5 branches; (10) seta 1-X inserted in a marginal notch or at the edge of the saddle; and (11) seta 14-P with 4–8 branches. Distinction of *An. cracens* based on pupal characters is more difficult, but the overall pigmentation of the pupa is distinctive in comparison to other members of the Dirus Complex. Especially evident is the lack of pigmentation of seta 9-III–VII, which are very light yellow, only slightly darker than the integument. Additionally for *An. cracens*, the male genital lobe of the pupa is noticeably constricted at middle length.

Anopheles (Cellia) scanloni Sallum and Peyton, sp.n.

Etymology: this species is named in honour of the late John E. Scanlon, whose studies on *Anopheles* represent an enormous contribution to knowledge of malaria epidemiology and vector species in Asia.

Anopheles dirus species C of Baimai *et al.*, 1984: 536 (cytogenetics); Wibowo *et al.*, 1984: 425 (H-banding pattern of sex chromosomes); Baimai *et al.*, 1987: 157 (crossing, chromosomes); May, 1987: 886 (neuroblast chromosomes); Baimai, 1988a: 13 (heterochromatin, evolution, distribution); Baimai, 1988b: 667 (population cytogenetics); Baimai *et al.*, 1988d: 372 (polytene chromosomes); Yasothornsrikul *et al.*, 1988: 703 (RFLP); Damrongphol & Baimai, 1989: 563 (egg morphology); Peyton, 1990: 197 (taxonomy); Sawadiparnich *et al.*, 1990: 477 (cytogenetics, crossing, evolution); Green *et al.*, 1992: 29 (enzyme electrophoresis); Audtho *et al.*, 1995: 107 (DNA hybridization); Poopittayasataporn & Baimai, 1995: 426 (polytene chromosomes, phylogeny); Walton *et al.*, 1999: 24 (ASA PCR species identification); Walton *et al.*, 2000: 962 (mtDNA genetic structure); Walton *et al.*, 2001: 569 (population genetics, microsatellites); Huong *et al.*, 2001: 615 (PCR species identification); Manguin *et al.*, 2002: 46 (SCAR markers, multiplex PCR).

Female. Head: proboscis length 1.64–2.27 mm (mean 1.95 mm), ratio of length to forefemur length 1.07–1.14 (mean 1.10), maxillary palpus length 1.52–2.15 mm (mean 1.82 mm), ratio of length to proboscis length 0.90–0.98 (mean 0.93), ratio of length to forefemur length 1.00–1.10 (mean 1.03), ratio of length of palpomeres 3/4 1.53–2.00 (mean 1.75), 3/5 2.18–2.71 (mean 2.43), 4/5 1.22–1.50 (mean 1.42), 4–5/3 0.91–1.08 (mean 0.99); palpomeres 2–4 with narrow apical white bands, pale scales of palpomere 5 white to pale cream-coloured, length of pale band of palpomere 5 1.00–3.50 (mean 1.87) length of basal dark band of palpomere 5, ratio of length of apical pale band of palpomere 4 to length of basal dark band of palpomere 5 0.28–2.00 (mean 0.79). Thorax: pleural setae as follows: 1,2 upper proepisternal, 0–2 prespiracular, 3–5 prealar, 3–5 upper mesokatepisternal, 2–4 lower mesokatepisternal, 2–4 upper mesepimeral. Wing: length 2.63–3.72 mm (mean 3.08 mm), pale scales on all veins light cream-coloured, spots on costa and subcosta slightly more obviously cream-coloured but not strongly contrasting with other spots, PHP spot of costa absent to prominent, HP spot reduced or small but rarely absent, PSP spot always present, reduced or prominent, SP spot always present, prominent, ASP spot always absent, PP spot 0.13–1.60 (mean 0.84) length of subcostal pale SCP spot, AD spot 1.10–4.75 (mean 2.10) length of PP spot, PSD spot of vein R extending basally from level 1 to level 5 on one or both wings, PSD spot of vein R entirely dark-scaled or with 1,2 pale interruptions, SD spot of vein R with 1–3 pale interruptions, 2–7 pale interruptions on PSD-PD spots of vein R on each wing, ratio of length of cell R_2 to R_{2+3} 1.25–1.70 (mean 1.39), ratio of length of cell R_2 to cell M_{1+2}

1.14–1.41 (mean 1.27). Legs: foretarsomeres 2 and 3 with complete or incomplete basal and apical pale bands, middle dark region of foretarsomere 2 entirely dark-scaled or with 1,3 pale spots, foretarsomere 4 with basal pale band, apical pale band reduced or absent, foretarsomere 5 mostly dark-scaled, apical pale scales reduced or absent, foretarsomeres 2–5 mostly pale-scaled along ventral surface, pale spots absent or less noticeable than on dorsal surface; midtarsomeres 2–4 usually dark-scaled with small apical patch of pale scales on dorsal surface, occasionally apical pale patches inconspicuous, midtarsomere 5 dark-scaled with faint apical pale band; hindtarsomeres 2, 3 mostly dark-scaled with small apical pale band, hindtarsomere 4 with apical and basal pale bands, hindtarsomere 5 with basal and apical pale bands, basal pale band sometimes absent or reduced to dorsal pale patch, apical pale band always present. Abdomen: tergum VI without scales, tergum VII with few sparse dark scales on posterior margin, tergum VIII with large posteromedial patch of yellowish to golden scales; sternum VI without scales, sternum VII with posteromedial patch of dark scales, sternum VIII variable, without scales or with few lateral pale scales, with anterolateral patches of pale scales.

Male. Essentially as in female except for sexual characters. Wing generally paler with reduced scaling, pale spots usually longer than in female. Palpomere 2 with long patch of pale scales at middle extending laterally, apex of palpomere 2 bare; palpomere 3 with long dorsal patch of pale scales at middle, extending laterally, apex of palpomere 3 with broad basal band of pale scales covering dorsal, lateral and ventral surfaces except for narrow, ventrolateral spot of dark scales, palpomeres 4 and 5 mainly pale-scaled with narrow basal band of dark scales. Abdomen: sternum VIII covered with yellowish to golden scales.

Pupa (Fig. 3, Table 5). Position and development of setae as figured; range and modal number of branches in Table 5. Measurements and counts from 20 to 30 specimens. In general as described for *An. latens* except for the following characters. Sterna II–VII with narrow, dark band near anterior margin. Cephalothorax: 5,11-CT frequently triple (5, 2–5; 11, 1–4), 6–10,12-CT usually double (see Table 5 for ranges). Abdomen: seta 6-I usually double (1–3), 7-I with 3–5 branches; 1-II with more than 20 fine branches, 6-II usually double (1,2), 7-II with 2–5 branches, 8,10,11-II absent, 9-II length 0.01–0.02 mm (mean 0.01 mm); 5-III with 6–10 branches, 9-III length 0.02–0.03 mm (mean 0.02 mm); 1,5-IV with 4–7 branches, 6-IV single, 9-IV length 0.03–0.10 mm (mean 0.05 mm), ratio of length of 9-IV/9-III 1.09–4.75 (mean 2.31), ratio of 9-IV/9-V 0.28–0.88 (mean 0.46); 1-V usually double (1–3), 5-V with 3–6 branches, 9-V usually simple, rarely with 1,2 minute denticles, length 0.08–0.13 mm (mean 0.11 mm); 1-VI usually double (1–3), 5-VI with 3–6 branches, 6-VI single or double, 9-VI usually simple, rarely with 1,2 minute denticles, length 0.10–0.13 mm (mean 0.12 mm); 1,6-VII single or double, 5-VII with 3–6 branches, 9-VII usually simple, rarely with 1, 2 minute denticles, length 0.11–0.14 mm (mean 0.12 mm); 9-VIII with 9–17 branches. Paddle: filamentous spicules on outer

apical margin either absent or sparse, always absent on inner margin; seta 2-Pa single; toothed margin index 0.82–0.90 (mean 0.86), paddle teeth narrow, tapering to acute apex.

Larva (Fig. 3, Table 6). Position and development of setae as figured; range and modal number of branches in Table 6. Measurements and counts from 18 to 40 specimens. In general as described for *An. latens* except for the following characters. Head: integument light brown to yellowish, with or without pattern of darkened spots, if present spots as follow: 1 small spot posteriorly at dorsal apotome and 2 well developed dark spots slightly anterior to it; a more developed, single dark spot posterior to seta 6-C in area between seta 8-C, and 2 small dark spots laterally; 2 lateral dark spots at dorsal apotome, posterior to seta 7-C; 1 dark spot ventrally on lateralia in the area of insertion of seta 12-C; a faint dark spot on lateralia bordering frontal ecdysial line; 1 dark spot on labiogula laterally at level of posterior tentorial pit; head length 0.60–0.73 mm (mean 0.67 mm), width 0.68–0.78 mm (mean 0.74 mm); antenna length 0.29–0.32 mm (mean 0.30 mm), ratio of distance between base and seta 1-A to antenna length 0.26–0.48 (mean 0.34); seta 2-C simple; 3-C length 0.07–0.10 mm (mean 0.09 mm); distance between bases of 2-C and 3-C 0.03–0.04 mm (mean 0.04 mm); 4-C single to triple, length 0.10–0.22 mm (mean 0.13 mm), extending beyond base of 2-C, distance between insertions of 2-C and 4-C 0.06–0.10 mm (mean 0.08 mm), ratio of length of 4-C to distance between insertions of 2-C and 4-C 1.20–3.26 (mean 1.70); distance between bases of 3-C and 4-C 0.05–0.08 mm (mean 0.07 mm); 5-C shorter than antenna, not reaching anterior margin of head, with 8–11 branches, 6-C with 10–14 branches, 7-C with 8–16 branches. Thorax: seta 1-P with 10–13 branches, tubercle of 1-P frequently separate from tubercle of 2,3-P, or rarely joined to tubercles of 2,3-P on one side, each tubercle with well developed apical tooth arising from posterodorsal side and projecting forward over bases of 1,2-P; 4-M with 2,3 branches, 6-M triple, 14-M with 7,8 branches; 3-T moderately developed, palmate, with 3,4 weak, nearly transparent lanceolate leaflets arising from a long narrow stem. Abdomen: seta 2-I usually double (1–3), 3-I single or double, 9-I with 3,4 branches; 1-II more developed than 1-I, with 10–16 leaflets, stem more pigmented than leaflets; 3-IV single or double, 13-IV triple, ratio of length of 13-IV to 10-IV 0.70–0.88 (mean 0.80); 1-VII with 9–13 leaflets, leaflets normally without apicolateral serrations, rarely few leaflets with minute apicolateral serrations; pecten teeth 12–17 with 4–6 long teeth alternating with 7–11 short teeth.

Material examined. HOLOTYPE: adult female with associated larval and pupal exuviae on microscope slide, acc. no. 08126–15, collected by AFRIMS team, 18 November 1979, deposited in the NMNH. Type locality: THAILAND, Kanchanaburi, Sai Yok (Office of Livestock) 14°21'N 98°59'E. PARATYPES: Kanchanaburi, Nam Tok, Khao Na Chang (14°13'N 99°5'E), 6 June 1965, acc. no. 0225 1M, 1F, 1Le, 1Pe; Phao Phu Thong (13°54'N 99°24'E), 21 September 1973, acc. no. 06537, 2F, 1Le, 1Pe; Huai Mae, Khamin (14°38'N 99°00'E), 24 November 1979, acc. no.




Fig. 3. *Anopheles scanloni* sp.n. Pupa (female): CT, cephalothorax; Pa, paddle; 1-IX, abdominal segments of pupa (dorsal on left, ventral on right), numbers on the left side denote seta 9 of designated segments. Larva: C, head; P, prothorax; M, mesothorax; T, metathorax; PP, pecten plate; abdominal segments I-VI left side dorsal, right side ventral; VII-X lateral (left side) view. Circle shows basal details of setae 1-3-P. Millimetre scale bars.

Table 5. Number of branches for pupal setae of *An. scanloni*: range (mode). n.c. = not counted.

Seta no.	Cephalothorax CT	Abdominal segments									Paddle P
		I	II	III	IV	V	VI	VII	VIII	IX	
0	–	–	1	1	1	1	1	1	1	–	–
1	2,3 (2)	n.c.	>20	5–10 (6)	4–7 (5)	1–3 (2)	1–3 (2)	1,2 (1)	–	1	1
2	2	3–6 (5)	4–7 (5)	3–6 (5)	2,3 (3)	2–4 (3)	2–4 (3)	2,3 (2)	–	–	1
3	2,3 (2)	1	1–3 (2)	1–3 (2)	5–8 (6)	2,3 (2)	1–3 (2)	1–3 (2)	–	–	–
4	1–4 (2)	2–6 (4)	3–7 (4)	1–5 (4)	2–5 (3)	3–5 (4)	1,2 (2)	1,2 (2)	1,2 (1)	–	–
5	2–5 (3)	1–3 (2)	2–7 (3)	6–10 (7)	4–7 (5)	3–6 (5)	3–6 (4)	3–6 (4)	–	–	–
6	1,2 (2)	1–3 (2)	1,2 (2)	1,2 (2)	1	1	1,2 (1)	1,2 (1)	–	–	–
7	2–4 (2)	3–5 (3)	2–5 (3)	1–4 (2)	1–4 (2)	1–3 (2)	1	1	–	–	–
8	1–3 (2)	–	–	1–3 (2)	1–3 (2)	1,2 (1)	1,2 (1)	1,2 (1)	–	–	–
9	1–3 (2)	1	1	1	1	1	1	1	9–17 (10)	–	–
10	1–4 (2)	–	1,2 (1)	1–4 (2)	1–3 (1)	1,2 (1)	1,2 (2)	1,2 (2)	–	–	–
11	1–4 (3)	–	–	1	1	1	1	1	–	–	–
12	1–3 (2)	–	–	–	–	–	–	–	–	–	–
13	–	–	–	–	–	–	–	–	–	–	–
14	–	–	–	–	1	1	1	1	1	–	–

Table 6. Number of branches for setae of fourth-instar larva of *An. scanloni*: range (mode). n.c. = not counted.

Seta no.	Head C	Thorax			Abdominal segments									
		P	M	T	I	II	III	IV	V	VI	VII	VIII	X	
0	1	1	—	—	—	1	1	1	1	1	1	1	—	
1	1	10–13 (13)	25–30 (26)	1	3–7 (5)	10–16 (10)	18–21 (19)	18–21 (19)	17–20 (17)	15–18 (17)	9–13 (12)	1	1	
2	1	7–12 (11)	2–3 (3)	1	1–3 (2)	5–7 (5)	5–6 (6)	3–5 (4)	3–5 (4)	4–5 (5)	4–6 (6)	5–6 (6)	13–14 (13)	
3	1	1	1	3–4 (4)	1–2 (1)	1	1	1–2 (2)	1	1	3	4–5 (5)	7	
4	1–4 (1)	13–18	2–3 (2)	3–4 (3)	4–6 (6)	4–6 (5)	2–3 (3)	2–3 (3)	2–3 (2)	1	1	1	9	
5	8–11 (9)	n.c.	1	28–37 (28)	3	4–7 (5)	3–4 (3)	3–5 (3)	3–5 (5)	4–5 (5)	4–5 (5)	3	—	
6	10–14 (11)	1	3	2	18–22 (20)	22–27 (22)	12–18 (12)	1–2 (2)	1–2 (2)	2–3 (3)	4–5 (4)	1-S	4–5 (4)	
7	8–16 (10)	14–25 (22)	2–4 (3)	27–33	14–17 (15)	19–22 (20)	4–5 (5)	4–6 (5)	4–6 (6)	3–5 (3)	3	2-S	5–6 (6)	
8	1–2 (2)	23–29 (27)	17–29 (23)	29–33	—	2	2	2–3 (2)	2–3 (3)	2–3 (3)	4–5 (5)	6-S	2	
9	3–5 (3)	1	1	1	3–4 (3)	6–8 (8)	7–8 (8)	6–7 (7)	6	5–8 (5)	4	7-S	1	
10	1–2 (2)	1	1	1	1	2	1	1	1	3–4 (3)	5–6 (5)	8-S	1–3 (2)	
11	26–29 (29)	1	1	1	2–3 (3)	1	2–3 (2)	2–3 (2)	2–3 (3)	2	2	9-S	1–3 (3)	
12	2	1–3 (1)	1	2–3 (2)	3–4 (3)	1	2–3 (2)	2	2	1	1	—	—	
13	5–8 (6)	4–5 (5)	4–5 (5)	2–3 (3)	4–5 (4)	3–6 (5)	4–7 (6)	3	2–3 (3)	5–9 (6)	2–3 (3)	—	—	
14	n.c.	5–8 (6)	7–8 (8)	—	—	—	1	1	1	1	1	1	—	
15	n.c	—	—	—	—	—	—	—	—	—	—	—	—	

08153, 5F, 6M, 10Le, 10Pe; Ban Plai Huai, Kaeng Rieng (14°24'N 99°5'E), 7 June 1982, acc. no. 09137(1), progeny, 16F, 12M, 28Le, 28Pe; Ban Tab Tao, Mu 2 (14°24'N 99°9'E), 8 June 1982, acc. no. 09139 (14), progeny, 5F, 4M, 8Le, 8Pe; Ban Huai Duan (14°24'N 99°5'E), 10 June 1982, acc. no. 09142(2), progeny, 19F, 24M, 42Le, 42Pe; Tha Kradan, Ban Phu Taka, Mu 3 (14°27'N 99°5'E), 1 July 1981, progeny, acc. no. 09120 (1), 12F, 11M, 26Le, 22Pe. All collected by AFRIMS staff team. Phattalung, Tanod, Ban Lo Chang Kra (7°10'N 99°30'E), February 1985, acc. no. PT15, progeny, 3F, 2Le, 2Pe, 1I, collected by V. Baimai. All paratypes deposited in the NMNH. Total material of *An. scanloni* examined: 1358 specimens comprising 221F, 228M, 431Le, 437Pe, 41I, derived from 18 separate collections from natural habitats (10 adult, 8 immature) and 28 progeny broods.

Distribution. *Anopheles scanloni* is known only from Thailand (Kanchanaburi, Nakhon Si Thammarat and Phattalung).

Bionomics. *Anopheles scanloni* may be an efficient vector of malaria parasites in South-east Asia (Manguin *et al.*, 2002). However, the precise involvement of the species in malaria transmission is not fully understood because morphological identification is problematical and *An. scanloni* has been confused with other members of the Dirus Complex. Adult females were obtained from human bait collections in evergreen forest, deciduous forest, secondary deciduous forest, cultivated field, corn plantation surrounded by forest, villages and 10 m from human habitation. Immatures of *An. scanloni* were taken from rock pools, ground pools and seepage springs. The water was always fresh, stagnant, coloured or clear, sometimes with some degree of pollution, and in partial shade. Larval habitats were in primary bamboo groves, tropical rain forest, deciduous forest and scrub environments situated in mountains and also in valleys at elevations from 75 to 480 m above sea level. Immatures were found in association with *An. barbirostris*, *An. kochi*, *An. maculatus* Theobald, *An. nigerrimus* Giles, *An. vagus* Dönitz, *Cx. fuscocephalus* Theobald, *Cx. infantulus* Edwards, *Cx. minor* (Leicester), *Oc. assamensis* (Theobald) and *Oc. gubernatoris* (Giles).

Systematics. Based on the distinctive H-banding pattern of metaphase sex chromosomes, Wibowo *et al.* (1984) distinguished *Anopheles scanloni* (as *An. dirus* C) from *An. balabacensis*, *An. dirus* and *An. cracens* (as *An. dirus* B). Several crossing, chromosomal and RFLP studies confirmed the species status of *An. scanloni* (Baimai *et al.*, 1984, 1987, 1988d; May, 1987; Baimai, 1988a, 1988b; Yasothornsrikul *et al.*, 1988; Sawadipaniich *et al.*, 1990; Audtho *et al.*, 1995; Poopittayasataporn & Baimai, 1995). Scanning electron micrographs of the eggs of *An. scanloni* and *An. dirus* showed that the pattern of the outer chorionic cells between the frill and the floats may be useful for distinguishing between these species (Damrongphol & Baimai, 1989). The banding pattern of the larval salivary gland polytene chromosomes showed interspecific differences,

particularly at the free ends of the X, 2R and 2L arms, between *An. scanloni*, *An. dirus* and *An. cracens* (Baimai *et al.*, 1988d). The species status of *An. scanloni* was also based on isozyme electromorphs (Green *et al.*, 1992), a non-radioactive DNA hybridization method (Audtho *et al.*, 1995), an allele-specific polymerase chain reaction of the ITS2 region of rDNA (Walton *et al.*, 1999), and nucleotide sequence data of the COI mitochondrial gene (Walton *et al.*, 2000). More recently, a PCR test employing species-specific primers was developed to separate four species of the Dirus Complex, including *An. scanloni* (Huong *et al.*, 2001), and a multiplex PCR-based method using species-specific primers (Manguin *et al.*, 2002). Studies on the population genetic structure of *An. scanloni* employing 11 microsatellite loci showed that there is a high degree of differentiation between the northern and the southern populations, suggesting the presence of two incipient species (Walton *et al.*, 2001).

The adult of *An. scanloni* can be recognized by the following combination of characters: (1) proboscis dark-scaled, slightly longer than forefemur, ratio of proboscis length to forefemur length 1.07–1.14 (mean 1.10); (2) PSD spot on vein R normally with 1, 2 pale interruptions; (3) ASP spot absent on costa; (4) hindtarsomere 4 with basal pale band; (5) apical white band on hindtibia with dark linear extension into basal portion of ventral surface (occurs only in specimens from Kanchanaburi) or without dark extension; (6) longest pale spot of vein 1A is at the level of the PSD spot; (7) PSD spot of vein R extending from level 1 to level 5 on one or both wings, most frequently to level 3 or 4; (8) pale spots and bands along dorsolateral surface of foretarsomeres 1 and 2 usually small, discrete, occupying less surface than the dark portions along this surface; (9) pale spots on foretarsomere 1 usually not fused, not forming long lines or splashes of pale scales along entire length, often restricted to 2–4 spots; (10) middle dark area of foretarsomere 2 usually without pale spot on dorsal surface, occasionally with 1–3 small pale spots.

The fourth-instar larva of *An. scanloni* can be recognized by the following combination of characters: (1) 3-C single; (2) seta 1-II not fully developed; (3) 5-C shorter than antenna, not reaching anterior margin of head; (4) seta 1-X inserted on the saddle; (5) tubercle of seta 1-P usually separate, occasionally narrowly joined to tubercle of 2,3-P on one side; (6) 2-C single and simple; (7) 4-C long, extending well beyond base of 2-C, ratio of length of 4-C to distance between the insertions of 2-C and 4-C 1.20–3.26; (8) length of seta 4-C varying from 0.10 to 0.22 mm (mean 0.13 mm).

The pupa of *An. scanloni* can be distinguished by the absence of a fringe of filamentous spicules on the inner margin of the paddle. However, *An. elegans* exhibits polymorphism for this character and some individuals may be misidentified as *An. scanloni*. Therefore, since *An. scanloni* and *An. elegans* are allopatric, geographical distribution should also be considered when separating *An. scanloni* from *An. elegans* on the basis of pupal characters alone.

Anopheles (Cellia) baimai Sallum & Peyton, sp.n.

Etymology: this species is named after Prof. Visut Baimai for his valuable contributions to systematics of the *Leucosphyrus* Group.

Anopheles dirus D of Baimai *et al.*, 1984: 536 (cytogenetics); Wibowo *et al.*, 1984: 425 (H-banding pattern of sex chromosomes); Baimai *et al.*, 1987: 157 (crossing, chromosomes); Baimai, 1988a: 13 (heterochromatin, evolution, distribution); Baimai, 1988b: 667 (population cytogenetics); Baimai *et al.*, 1988d: 372 (cytological differences, chromosome rearrangement); Baimai *et al.*, 1988e: 661 (chromosomal polymorphism, distribution); Panyim *et al.*, 1988: 47 (DNA probe); Yasothornsrikul *et al.*, 1988: 703 (RFLP); Damrongphol & Baimai, 1989: 563 (egg morphology); Peyton, 1990: 197 (taxonomy); Green *et al.*, 1992: 29 (enzyme electromorphs); Audtho *et al.*, 1995: 107 (DNA hybridization); Poopittayasataporn & Baimai, 1995: 426 (polytene chromosomes, phylogeny); Rattananarithikul *et al.*, 1995: 428 (bionomics); Xu & Qu, 1997: 134 (ITS2 marker); Xu *et al.*, 1998: 385 (ITS2 marker); Walton *et al.*, 1999: 24 (ASA PCR species identification); Walton *et al.*, 2000: 962 (mtDNA genetic structure); Huong *et al.*, 2001: 615 (PCR species identification); Walton *et al.*, 2001: 569 (genetics, microsatellites); Manguin *et al.*, 2002: 46 (SCAR multiplex PCR).

Anopheles dirus of Rosenberg, 1982: 192 (breeding habits); Rosenberg & Maheswary, 1982: 183 (bionomics); Dutta *et al.*, 1989a: 149 (sporozoite detection); Dutta *et al.*, 1989b: 95 (bionomics, sporozoite detection); Dutta *et al.*, 1991: 121 (vector incrimination); Dutta *et al.*, 1992: 245 (sporozoite detection); Dutta *et al.*, 1993: 135 (bionomics); Misra *et al.*, 1993: 62 (bionomics); Nandi *et al.*, 1993: 18 (sporozoite infection); Dutta *et al.*, 1996: 378 (feeding behaviour); Nandi *et al.*, 1996: 279 (bionomics); Dutta *et al.*, 1997: 204 (bionomics); Prakash *et al.*, 1997a: 31 (biting behaviour, parity rate); Prakash *et al.*, 1997b: 117 (bionomics); Prakash *et al.*, 1997c: 610 (bionomics, feeding behaviour); Khan *et al.*, 1998: 131 (bionomics); Prakash *et al.*, 1998a: 291 (insecticide susceptibility); Prakash *et al.*, 1998b: 191 (bionomics); Moore *et al.*, 2000: 282 (bionomics); Nandi *et al.*, 2000: 95 (feeding habits); Prakash *et al.*, 2000a: 97 (bionomics); Prakash *et al.*, 2000b: 121 (bionomics); Prakash *et al.*, 2001: 231 (vectorial capacity); Prakash *et al.*, 2002: 95 (breeding habitats).

Female. Head: proboscis length 1.97–2.38 mm (mean 2.09 mm), ratio of length to forefemur length 1.01–1.14 (mean 1.08), maxillary palpus length 1.77–2.18 mm (mean 1.96 mm), ratio of length to proboscis length 0.88–1.00 (mean 0.94), ratio of length to forefemur length 0.96–1.05 (mean 1.00), ratio of length of palpomeres 3/4 1.50–1.67 (mean 1.63), 3/5 2.08–2.78 (mean 2.27), 4/5 1.25–1.67 (mean 1.39), 4–5/3 0.96–1.20 (mean 1.06), palpomeres 2–4 with narrow apical silvery white bands, pale scales of palpomere 5 white to pale cream-coloured, length of pale band of palpomere 5 1.00–2.33 (mean 1.39) length of basal dark band of palpomere 5, ratio of length of apical pale band of palpomere 4 to length of basal dark band of palpomere 5 0.37–1.00 (mean 0.64). Thorax: pleural setae as follow: 1–3

upper proepisternal, 0–4 prespiracular, 2–6 prealar, 3–5 upper mesokatepisternal, 1–3 lower mesokatepisternal, 3,4 upper mesepimeral. Wing: length 3.09–3.54 mm (mean 3.39 mm), pale scales of all veins light cream-coloured, spots on costa and subcosta more obviously cream-coloured, sometimes bordering on pale golden, pale spots on remaining veins lighter, not strongly contrasting with others; ASP spot always absent, PP spot 0.38–2.00 (mean 1.01) length of SCP spot, AD spot 1.15–4.17 (mean 2.11) length of PP spot, PSD spot of vein R extending basally from level 1 to level 5 on one or both wings, PSD spot of vein R entirely dark or with 1–4 small pale spots, sum of pale spots on PSD-PD spots of vein R 3–10 for each wing; ratio of length of cell R_2 to vein R_{2+3} 1.17–1.77 (mean 1.48), ratio of length of cell R_2 to cell M_{1+2} 1.12–1.26 (mean 1.19). Legs: foretarsomere 2 variable, sometimes entirely pale-scaled along dorsal and lateral surfaces, or dark-scaled with basal and apical pale bands, middle dark area variable, usually mostly pale along dorsal and lateral surfaces, dark scales restricted to 1, 2 middle dark bands, less frequently mostly dark-scaled, with or without 1, 2 pale spots, foretarsomere 3 dark-scaled with basal and apical pale bands, foretarsomere 4 usually dark-scaled with basal and apical pale patches, occasionally dark part less distinct, pale scales varying from light brown to tan, foretarsomere 5 entirely pale-scaled or with indistinct tan to light brown basal band, pale at apex, foretarsomeres 2–5 tan-scaled along ventral surface, basal and apical pale bands less noticeable or absent on ventral surface; midtarsomeres 2–4 with apical patches of pale scales more evident on dorsal surface, sometimes pale patches poorly evident, midtarsomere 2 variable, entirely dark-scaled or with 1, 2 pale spots on dorsal surface, midtarsomere 5 variable from tan to light brown or dark at base and pale at apex; hindtarsomeres 2 and 3 with apical pale bands, hindtarsomere 2 with 1–3 pale spots on middle dark region, occasionally reduced to 1, 2 pale scales, hindtarsomeres 4 and 5 with basal and apical pale bands, sometimes reduced to few basal and apical pale scales on hindtarsomere 5. Abdomen: tergum VI sometimes with 2–4 pale cream-coloured scales on posteromedial margin, tergum VII with few brownish scales on posterior margin, tergum VIII covered with cream-coloured to yellowish to golden scales, scales sometimes reduced to small posteromedial patch; sternum VI usually without scales or occasionally with 1–4 dark posteromedial scales.

Male. Essentially as in female except for sexual characters. Wing generally paler with reduced scaling, pale spots usually longer than in female. Palpomere 2 with dorsal patch of pale scales at middle, apex of palpomere 2 bare; palpomere 3 with long dorsal patch of pale scales at middle extending laterally, apex of palpomere 3 with broad band of pale scales covering dorsal, lateral and ventral surfaces except for narrow ventrolateral line of dark scales at apex; palpomeres 4 and 5 pale-scaled with basal band of dark scales, sometimes reduced to very narrow dark band on palpomere 5, more visible on dorsal and ventral surfaces, scales on lateral surface entirely pale or tan at base, dark band of palpomere 5 with ventral dark extension, not reaching apex of palpomere. Abdomen: sternum VIII covered

with white to pale cream-coloured scales centrally, scales yellowish laterally.

Pupa (Fig. 4, Table 7). Position and development of setae as figured; range and modal number of branches in Table 7. Measurements and counts from 20 to 38 specimens. In general as described for *An. latens* except for the following characters. Sterna II-VII with narrow dark bands near anterior margins, usually darker on segments II-V. Cephalothorax: seta 5-CT with 3-7 branches, 6,7,8-CT usually double (6,8, 1-3; 7, 2 or 3), 10-CT with 2-5 branches. Seta 9-IV-VIII light to medium-brown, slightly darker at base. Abdomen: seta 6-I usually double (2,3), 7-I with 3-5 branches; 6-II single to 5-branched, 7-II with 2-5 branches, 8,10,11-II absent, 9-II length 0.01-0.02 mm (mean 0.01 mm); 5-III with 6-14 branches, 9-III length 0.01-0.02 mm (mean 0.02 mm); 5-IV with 4-10 branches, 6-IV single or double, 9-IV length 0.02-0.09 mm (mean 0.04 mm), ratios of length of seta 9-IV/9-III 1.15-4.40 (mean 2.15) and seta 9-IV/9-V 0.20-0.84 (mean 0.42); 5-V with 4-7 branches, 6-V always single, 9-V length 0.04-0.12 mm (mean 0.10 mm); 1-VI usually double (1-3), 6-VI single, 9-VI length 0.07-0.13 mm (mean 0.10 mm); 1-VII single or double, 5-VII with 3-7 branches, 9-VII length 0.09-0.14 mm (mean 0.11 mm); 9-VIII with 9-16 branches. Paddle: 2-Pa single or double; toothed margin index 0.76-0.88 (mean 0.84), paddle teeth well developed, teeth tapering to acute apex, teeth widely separated.

Larva (Fig. 4, Table 8). Position and development of setae as figured; range and modal number of branches in Table 8. Measurements and counts from 15 to 40 specimens. In general similar to *An. latens* except for the following characters. Head: integument light brown to yellowish, usually without pattern of dark spots but if present as follows: 2 small dark spots posteriorly on dorsal apotome; one dark spot slightly anterior them; one dark spot centrally on dorsal apotome posterior to seta 5-C; 2 lateral spots on dorsal apotome between insertions of setae 7-C and 8-C along dorsal ecdysial line; 2 dark spots on lateralialia, 1 placed in area of 12-P insertion and 1 slightly posterior; and 2 dark spots on labiogula at level of hypostomal suture; head length 0.62-0.75 mm (mean 0.67 mm), width 0.66-0.81 mm (mean 0.76 mm); antenna length 0.28-0.34 mm (mean 0.31 mm), ratio of distance between base and seta 1-A to antenna length 0.25-0.40 (mean 0.32); seta 2-C usually simple, rarely with 1, 2 minute spicules on apical 0.5, 3-C length 0.06-0.10 mm (mean 0.08 mm), distance between bases of 2-C and 3-C 0.03-0.04 mm (mean 0.04 mm), 4-C posterolaterad of 2-C, single or double, length 0.07-0.12 mm (mean 0.09 mm), extending to or beyond base of 2-C, distance between insertions of 2-C and 4-C 0.08-0.13 mm (mean 0.10 mm), ratio of length of 4-C to distance between insertions of 2-C and 4-C 0.67-1.31 (mean 0.94), distance between bases of 3-C and 4-C 0.08-0.13 mm (mean 0.09 mm), 5-C longer than antenna, not reaching anterior margin of head, with 8-12 branches, 6-C with 10-16 branches, 7-C with 9-14 branches. Thorax: tubercles of all large setae yellowish to light brown; seta 1-P with 13-22 branches, tubercle of 1-P joined to tubercle of 2,3-C, each tubercle with prominent, apical tooth arising

from posterodorsal side and projecting forward over bases of each seta, 14-P with 4-7 branches; 4-M with 2,3 branches, 14-M with 4-8 branches; 3-T moderately developed, somewhat palmate, with 3-8 leaflets arising from long slender stem. Abdomen: seta 1-I small, somewhat palmate, with 4-8 leaflets, 2-I single to triple; 1-II similar to 1-I with 4-13 leaflets; 3-IV double or triple, 13-IV with 3,4 branches, ratio of length of seta 13-IV to 10-IV 0.46-0.88 (mean 0.71); 1-VII with 10-16 lanceolate leaflets, leaflets frequently smooth, sometimes with minute apicolateral serrations, apical filament poorly defined; 1-X long, single, inserted in marginal notch or on edge of saddle; 11-16 pecten teeth, with 3-5 long teeth alternating with 8-11 short teeth.

Material examined. HOLOTYPE: adult female with associated larval and pupal exuviae on microscope slide, acc. no. TH1690(11)-15, collected by V. Baimai, 7 August 1989, progeny, deposited in the NMNH. Type locality: THAILAND, Mae Sot, Ban Kariang, Thum Rua (16°40'N 98°40'E). PARATYPES: BANGLADESH, Chittagong, Sylhet, Chaklapungee Tea Estate (24°00'N 91°25'E), 27 June 1975, acc. no. [072], 11F, 12M, 19Le, 17Pe; 21 November 1975, acc. no. [372], 1F, 1M, 3Le, 3Pe, 4I; 28 December 1975, acc. no. [412], 1F, 2Le, 2Pe, 2I, all collected by R. Rosenberg. THAILAND, Chumpton, Wang Mai, Ban Jo Po Ro (10°29'N 98°55'E), 9 September 1978, acc. no. 07970(4), progeny, 7F, 3M, 9Le, 10Pe. Kanchanaburi, Huai Mae Nam Noi (14°24'N 98°49'E), 28 May 1965, acc. no. 0161, 7F, 5M, 6Le, 10Pe, 2I; Sangkhla Buri, Ban Wang Kalang (15°6'N 98°28'E), 4 July 1974, acc. no. 06865, 3F, 2Le, 2Pe (all collected by AFRIMS staff team). Krabi, Plai Phraya, Ban Bang Haeng (8°34'N 98°44'E), November 1986, acc. no. KB50, 3F, 3M, 5Le, 6Pe; acc. no. KB106, 1F, 1M, 2Le, 2Pe (collected by V. Baimai). Mae Hong Son: Mae Sariang, Sop Muei (17°53'N 97°58'E), 2-3 October 1985, acc. no. TH(5), progeny, 4F, 10M, 7Le, 11Pe, 4I; Ban Mae Ho Nua (18°8'N 98°3'E), 15 September 1966, acc. no. 01523, 1F, 1Pe; acc. no. 01615, 6F, 2M, 2Le, 7Pe, 4 L (all collected by AFRIMS team). Nakhon Si Thammarat, Tung Yai, Ban Tham Phae Dan (8°10'N 99°20'E), 3-8 December 1985, acc. no. TY58, progeny, 6F, 4M, 10Le, 10Pe. Phangnga: Khao Pak Chaung (Chong) (8°17'N 98°23'E), 13 October 1966, acc. no. 01650, 4F, 1M, 1Le, 4Pe; Ban Bang Kaeo (8°35'N 98°32'E), 23 February 1986, acc. no. PG-30(F-3), colony, 8F, 1M, 8Le, 8Pe (all collected by V. Baimai). Tak: Tha Song Yang, Mae Tan, Komali Ko (17°12'N 98°14'E), 5-8 July 1987, acc. no. TH509(1), 3F, 8M, 11Le, 11Pe, 5I; Mae Sot, Ban Kariang, Thum Rua (16°40'N 98°40'E), 7 August 1989, acc. no. TH1690(10), 11F, 3M, 12Le, 12Pe, 2 L (all collected by V. Baimai). All paratypes deposited in the NMNH. Total material of *An. baimaii* examined: 5620 specimens comprising 1355F, 898M, 1404Le, 1657Pe, 306I, derived from 295 separate collections from natural habitats (74 adult, 225 immature) and 84 progeny broods.

Distribution. *Anopheles baimaii* is known from Bangladesh (Chittagong), India (Andaman Islands, Assam, Meghalaya,




Fig. 4. *Anopheles baimai* sp.n. Pupa (female): CT, cephalothorax; Pa, paddle; I-IX, abdominal segments of pupa (dorsal on left, ventral on right), numbers on the left side denote seta 9 of designated segments. Larva: C, head; P, prothorax; M, mesothorax; T, metathorax; PP, pecten plate; abdominal segments I-VI left side dorsal, right side ventral; VII-X lateral (left side) view. Circle shows basal details of setae 1-3-P. Millimetre scale bars.

Table 7. Number of branches for pupal setae of *An. baimaii*: range (mode). n.c. = not counted.

Seta no.	Cephalothorax CT	Abdominal segments									Paddle P
		I	II	III	IV	V	VI	VII	VIII	IX	
0	-	-	1	1	1	1	1	1	1	-	-
1	2-3 (3)	n.c.	>30	5-10 (7)	3-7 (5)	2-4 (3)	1-3 (2)	1-2 (1)	-	1	1
2	2-4 (2)	4-9 (5)	4-8 (5)	4-7 (6)	3-5 (3)	2-5 (3)	2-4 (3)	2-4 (2)	-	-	1-2 (2)
3	2-4 (2)	1	2-4 (3)	2-4 (3)	5-8 (7)	2-4 (2)	1-2 (2)	2-6 (3)	-	-	-
4	2-3 (2)	4-9 (6)	4-7 (5)	4-8 (4)	2-7 (5)	3-7 (5)	1-3 (2)	1-3 (2)	1-3 (2)	-	-
5	3-7 (4)	2-4 (3)	2-5 (3)	6-14 (8)	4-10 (5)	4-7 (7)	3-7 (5)	3-7 (6)	-	-	-
6	1-3 (2)	2-3 (2)	1-5 (2)	2-3 (2)	1-2 (1)	1	1	1-3 (2)	-	-	-
7	2-3 (2)	3-5 (3)	2-5 (3)	2-5 (3)	2-5 (3)	2-5 (3)	1-2 (1)	1-2 (1)	-	-	-
8	1-3 (2)	-	-	2-5 (3)	2-4 (2)	2-3 (3)	1-3 (2)	2-4 (3)	-	-	-
9	2-3 (3)	1	1	1	1	1	1	1	9-16 (13)	-	-
10	2-5 (3)	-	-	2-4 (3)	1-2 (1)	1-2 (2)	1-3 (2)	2-4 (2)	-	-	-
11	3-5 (3)	-	-	1	1	1	1	1-3 (2)	-	-	-
12	2-4 (3)	-	-	-	-	-	-	-	-	-	-
13	-	-	-	-	-	-	-	-	-	-	-
14	-	-	-	-	1	1	1	1	1	-	-

Table 8. Number of branches for setae of fourth-instar larva of *An. baimaii*: range (mode). n.c. = not counted.

Seta no.	Head C	Thorax			Abdominal segments									
		P	M	T	I	II	III	IV	V	VI	VII	VIII	X	
0	1	1	—	—	—	1	1	1	1	1	1	1	—	
1	1	13–22 (16)	25–38 (28)	1	4–8 (6)	4–13 (8)	16–24 (19)	15–22 (16)	16–21 (17)	12–20 (17)	10–16 (13)	1	1	
2	1	9–15 (10)	1–5 (2)	1	1–3 (2)	5–8 (7)	5–7 (6)	3–5 (4)	3–5 (4)	4–5 (4)	4–7 (6)	4–8 (6)	13–23 (16)	
3	1	1	1	3–8 (5)	1–3 (1)	1–2 (1)	1	2–3 (3)	1	1	2–4 (2)	4–8 (6)	8–11 (10)	
4	1–2 (1)	12–20 (13)	2–3 (3)	3–5 (3)	4–7 (5)	4–9 (5)	2–4 (3)	3–4 (3)	3–4 (3)	1	1	1	9	
5	8–12 (9)	n.c.	1	23–40 (33)	3–4 (3)	4–6 (5)	3–4 (3)	3–4 (3)	3–5 (3)	3–5 (4)	3–6 (5)	3–5 (3)	—	
6	10–16 (14)	1	3–6 (3)	1–3 (3)	18–21 (20)	17–26 (20)	13–18 (15)	2–3 (2)	2–3 (2)	2–3 (3)	2–5 (4)	1-S	3–6 (5)	
7	9–14 (12)	18–25 (20)	2–3 (3)	21–29	15–19 (15)	15–26 (19)	3–6 (5)	4–7 (4)	3–6 (5)	3–4 (3)	2–4 (3)	2-S	4–7 (6)	
8	1–2 (1)	22–31 (27)	17–28 (26)	24–28 (24)	—	1–2 (2)	2	2–3 (2)	2–3 (2)	2–3 (2)	3–5 (4)	6-S	1–3 (2)	
9	1–4 (2)	1	1	1	3–5 (4)	5–12 (6)	7–9 (8)	6–9 (8)	4–7 (6)	4–8 (6)	3–6 (4)	7-S	1	
10	1–2 (1)	1	1	1	1	2–4 (2)	1	1	1	2–3 (3)	3–7 (5)	8-S	2–3 (3)	
11	24–30 (30)	1	1	1	3–4 (3)	1	2–4 (3)	2–4 (3)	2–4 (3)	2–4 (2)	1–4 (2)	9-S	2–4 (3)	
12	1–2 (1)	1	1	2–3 (2)	3–4 (3)	1	2–3 (2)	2–3 (3)	2–3 (3)	1	1	—	—	
13	4–7 (5)	1–7 (5)	4–6 (5)	2–3 (3)	3–5 (4)	4–6 (5)	3–7 (5)	3–4 (3)	3–4 (3)	3–8 (6)	2–4 (2)	—	—	
14	n.c.	4–7 (5)	4–8 (6)	—	—	—	—	1	1	1	1	1	—	
15	3–5 (5)	—	—	—	—	—	—	—	—	—	—	—	—	

West Bengal), Myanmar (Tenasserim, Pegu, Sagaing Divisions), Thailand (Chumphon, Kanchanaburi, Krabi, Mae Hong Son, Nakhon Si Thammarat, Phangnga, Ranong, Tak Provinces), and Yunnan Province, southern China.

Bionomics. Females of *An. baimaii* were collected from cowsheds and hospitals in Assam, Cachar District, India (Ramsay, 1930) and at a settlement of brick-makers in a series of constructions made of bamboo, called 'bashas', at Chunati near Chringa, Bangladesh (Macan, 1950). *Anopheles baimaii* seems to be an important vector of human malaria parasites in some areas in Thailand, Burma, Bangladesh and northeastern India (Rahman *et al.*, 1977; Rosenberg & Maheswary, 1982; Rosenberg, 1982; Dutta *et al.*, 1989a, 1989b, 1991; Green *et al.*, 1991; Prakash *et al.*, 1997b, 2001; Xu *et al.*, 1998; Chareonviriyaphap *et al.*, 2000). For example, *An. baimaii* was the vector responsible for holoendemic malaria in a forest community of Bangladesh, with up to 3.8% sporozoite rate, although transmission occurred only during the 7-month monsoon. Females were both exophagic and endophagic with biting activity pattern irregular, related neither with rainfall nor other climatic parameters. Generally the biting activity peaked from 22.00 to 02.00 hours. Also, mostly adult female lives long enough to reach infectivity to *P. falciparum* (Rosenberg & Maheswary, 1982). Numbers biting indoors were greatly diminished in DDT-sprayed houses, and they fed earlier than in unsprayed houses. In Assam, India, in an isolated village of the rainforest-fringe, during the monsoon season, the vectorial capacity of *An. baimaii* for *P. vivax* was higher than for *P. falciparum*, decreasing after the monsoon to zero in the cold dry season (Prakash *et al.*, 2001). Malaria prevalence and vector density were higher in localities closer to the forest in Naga Hills with *An. minimus* as well as *An. baimaii* as vectors (Prakash *et al.*, 2000b). In upper Assam, biting activity of *An. baimaii* began at 19.00 hours, ended at 05.00 hours and was most intense between 21.00 and 03.00 hours (Prakash *et al.*, 2005). During the post-monsoon season, peak biting activity was from 03.00 to 05.00 hours, whereas in the pre-monsoon and monsoon season it was from 21.00 hours to midnight. The anthropophilic index was 92.3% (Prakash *et al.*, 1997a). In north-eastern areas of India, *An. baimaii* is highly anthropophilic, biting throughout the night with several activity peaks from 20.00 to 03.00 hours (Dutta *et al.*, 1996). Light trap collections in human dwellings in villages at the fringe of forest in Dibrugarh District, Assam, showed that *An. baimaii* was one of the most common mosquito species frequenting houses but it was the only anopheline species collected in an isolated village (Prakash *et al.*, 1997b, 1998b). Species density was related to seasonality, peaking in July, but it was very low in the dry, cool months. Positive correlation was observed between mosquito density and rainfall that occurred 2 weeks prior to collections (Prakash *et al.*, 1997b). In Nagaland, India, *An. baimaii* was found in low densities, but was more abundant during the monsoon season (Misra *et al.*, 1993). *Anopheles baimaii* was the most common of 12 *Anopheles* species collected in Changland District, Arunachal Pradesh in July and August, and some individuals were found positive for sporozoites

in Tirap and Arunachal Pradesh Districts, Assam, where sporozoites were found in the salivary glands of one specimen (Dutta *et al.*, 1989a, b, 1992). During a longitudinal study of malaria transmission carried out in three villages in Boko PHC, Assam, at the same time that indoor DDT spraying was being used for malaria control, *An. baimaii* was found in low numbers, and few specimens were positive for sporozoites (Nandi *et al.*, 1993). The sporozoite rate was 1.6% and the parous rate was 64.7% in an isolated village at the forest fringe in Assam, where malaria transmission increased with the increase in density of *An. baimaii* (Prakash *et al.*, 1997b). In north-eastern localities in West Bengal, *An. baimaii* was observed to be an anthropophilic species (Nandi *et al.*, 2000). Also, in Jalpaiguri duars of west Bengal, *An. baimaii* was more common during rainy months and found in close association with human environments (Nandi *et al.*, 1996).

In Thailand, adults of *An. baimaii* were collected biting humans in primary and secondary evergreen forest, primary and secondary deciduous forest, primary and secondary rain forest, secondary rain forest mixed with fruit plantations, orchards and rubber plantations, banana and pineapple plantations, villages, villages with fruit plantations, inside houses, and also in the canopy of rain forests 16.5 m above ground level.

The larval habitats of *An. baimaii* are similar to those of *An. dirus*. Although *An. baimaii* can share the same habitats with *An. dirus*, the former is more common in temporary breeding places than the latter (Rattananarithikul *et al.*, 1995). Immatures of *An. baimaii* have been found in domestic elephant and animal footprints, wheels and cart tracks along paths and roads, animal wallows, seepage pools, flood pools, pools in drying and dried streambeds and stream margins, ground pools, flood pools, large flood pools, seepage pools, limestone pools, sandy pools in banks of streams, shallow streams, ditches, flooded ditches, charcoal pits, small and large rock pools, swamps and rice paddies. The water was always fresh, stagnant, clear, turbid or sometimes with some degree of pollution. The pools were generally less than 2 m wide and 30 cm deep, under partial or heavy shade, or sometimes in sunlit areas. The larval habitats were inside primary bamboo grove forest, primary and secondary rain forest, primary and secondary deciduous forest, primary and secondary evergreen forest, secondary scrub in villages, secondary forest mixed with fruit plantations, evergreen rain forest, rubber tree plantations, village with fruit plantations, and orchards and villages located in valleys, hilly and mountainous areas.

In India, immatures of *An. baimaii* were collected in a well, 100 m above sea level. In Upper Brahmaputra Valley, Assam, immatures were found exclusively in forested areas (Khan *et al.*, 1998), whereas in a forest-fringe village in Assam, immatures were taken from small, shallow, rain-filled, temporary, shady and partly shaded puddles and ground pools during rainy months. In dry months, immatures were taken from permanent streams in the forest adjoining a village (Prakash *et al.*, 1997c). In Changland District, Arunachal Pradesh, *An. baimaii* is also a forest species found in small stagnant ground pools, completely or partially shaded by plants in open jungle (Dutta *et al.*, 1989b).

In Bangladesh, during the annual 6-month monsoon season, immatures of *An. baimaii* was found in puddles on footpaths, and in turbulence pits at the heads of drainage gullies that held water for at least 5 days. In the dry season, the breeding places were in streams about 3 km distant from the malaria focus. Immatures seem to be well adapted to small temporary pools. Eggs were laid above the water line and remained viable for about 2 weeks. Interestingly, larvae of *An. baimaii* were observed to leave a draining pool and crawl as far as 53 cm, sometimes to another pool. Also, larvae remained alive up to 94 h at the bottom of drained pools (Rosenberg, 1982).

During field collections carried out for the present study, immatures were collected from turbulence pit ground pools, turbulence pits, a narrow sand gully head, large pools, large turbulent pools in the middle of a stream bed, small ground pools, ground pools in a foot path, ground pools under a large steel bridge, stream pools, stream bed pools, head waters of streams, stream margins, cave pools, pools along roads, culvert pools, riverside pools, ditches on roads, ditches in streams, puddles in villages, sluggish streams, a stream bed below factory tank seepage, tank runoff and a pothole. The water was always fresh, stagnant, clear, turbid, sometimes dark or brownish, usually in heavy or partial shade or in full sun. The larval habitats were situated in tropical rain forest, in mountainous, hilly areas or in valleys.

In Yunnan Province, southern China, larvae of *An. baimaii* were found in habitats situated near villages; however, no adults were captured in CDC light traps with UV or incandescent bulbs during 57 night collections (Moore et al., 2001). Baimai et al. (1988e) showed that *An. baimaii* is highly polymorphic for chromosome inversions and that the distribution of the polymorphisms may be associated with the geography and epidemiology of malaria transmission.

Systematics. Using evidence from the heterochromatin of the metaphase sex chromosomes, Baimai et al. (1984) found support for a fourth species within the Dirus Complex, which was designated as *An. dirus* D. The species status of *An. baimaii* was confirmed by: crossing experiments and chromosomal differences (Baimai et al., 1987, 1988e); a DNA probe based on 7000 recombinant clones derived from wild females (Panyim et al., 1988); a non-radioactive DNA probe method (Audtho et al., 1995); nucleotide sequence of the ITS2 of rDNA (Xu & Qu, 1997; Xu et al., 1998); allele-specific amplification in the nucleotide sequence of ITS2 (Walton et al., 1999); a PCR based method (Huong et al., 2001); and a SCAR marker and multiplex PCR-based method (Manguin et al., 2002). Additionally, Green et al. (1991) found support for the hypothesis that the four chromosomal/electrophoretic forms of the Dirus Complex represented four distinct biological species, and Damrongphol & Baimai (1989) pointed out diagnostic morphological characters of the egg, useful for distinguishing *An. baimaii* from the other members of the complex. The phylogenetic relationships among five species of the complex, including *An. baimaii*, were investigated employing data on the fixed inversion of

3Rb and inversion polymorphism of the X chromosome (Poopittayasataporn & Baimai, 1995). Using 11 microsatellite loci, Walton et al. (2001) showed that *An. dirus*, *An. baimaii* and *An. scanloni* are very distinctive; however, no differentiation was observed either within or between *An. dirus* and *An. baimaii* based on COI gene sequences. The genetic diversity was higher in *An. baimaii* than in *An. dirus*, suggesting that expansion occurred first in *An. baimaii* and then in *An. dirus* (Walton et al., 2000).

Adults of *An. baimaii* can be recognized by the following combination of characters: (1) proboscis dark-scaled, slightly longer than forefemur, ratio of proboscis length to forefemur length 1.01–1.14 (mean = 1.08); (2) PSD spot of vein R rarely entirely dark-scaled, frequently with 1 or 2 pale interruptions on one or both wings; (3) ASP spot absent on costa and usually on subcosta; (4) hindtarsomere 4 with pale scales dorsally at base; (5) hindtarsomere 5 with a minute basal pale band; (6) apical pale band on hindtibia entirely white, without a dark extension into the basal portion; (7) vein 1A occasionally with a long pale spot at level of the PSD spot at least on one wing, when present this spot is always the longest pale spot of vein 1A; (8) combined pale spots and bands along the dorsolateral surface of foretarsomeres 1 and 2 dominate the dark portions along this line, often two or more of these pale spots on tarsomere 1 are fused, forming long lines or splashes of pale scales; (9) foretarsomere 2 often has the pale band and spots fused, is completely pale dorsally, or the pale spots and bands are longer, reducing the dark areas in the middle to narrow bands; (10) the PSD spot on vein R usually extends basally from level 1 to level 5, most often to level 1, level 3 and level 4.

The fourth-instar larva of *An. baimaii* can be distinguished from other members of the Dirus Complex except *An. cracens* by the following combination of characters: (1) 5-C noticeably longer than antenna; (2) basal sclerotized tubercle of seta 1-P with a prominent tooth or spine arising from the posterodorsal margin; (3) seta 4-C short, usually not reaching or extending beyond the base of 2-C, less frequently extending 0.3 beyond base of 2-C, the ratio of 4-C length to the distance between the insertions of 2-C and 4-C 0.67–1.31 (mean = 0.94); (4) seta 3-C extends to or slightly beyond anterior margin of head; (5) seta 1-X inserted in a marginal notch or at the edge of the saddle; and (6) seta 14-P with 4–8 branches.

Except for *An. scanloni* and the male pupa of *An. cracens*, the pupal stage of *An. baimaii* is indistinguishable from those of the other members of the Dirus Complex. As mentioned above, the pupal paddle of *An. scanloni* does not have a fringe of spicules on its inner margin, whereas it is always present in *An. baimaii*. Additionally, in the pupa of *An. cracens*, the male genital lobe is noticeably constricted at midlength, but not constricted in *An. baimaii*.

Anopheles (Cellia) elegans (James)

Myzomyia (?) *elegans* James, 1903. In Theobald, 1903: 51 (F* wing, scales) (new species of James, possibly a variety of

An. leucosphyrus). Type specimens: Holotype pinned female, BMNH, London. Type locality: India, Karnataka, Kârwâr.

Anopheles balabacensis of Reid, 1970: 56 (*in part*, systematics).

Anopheles dirus species E of Tewari *et al.*, 1987: 25 (bionomics, distribution); Peyton & Ramalingam, 1988: 274; Bhat, 1988: 104 (distribution); Peyton, 1990: 197 (taxonomy); Sawadipani *et al.* 1990: 25 (cytogenetics, crossing).

Anopheles elegans of Townsend, 1990: 36 (type information).

Anopheles leucosphyrus of Christophers, 1933: 177 (type information).

Anopheles leucosphyrus balabacensis of Colless, 1956: 76 (taxonomic notes, type information).

Anopheles leucosphyrus var. *elegans* of Cogill, 1903: 330 (larval bionomics, *in part*); James & Liston, 1940: 82 (wing, page 83); Reid, 1949: 46 (taxonomic notes, *in part*, type information).

Neomyzomyia elegans of Theobald, 1910: 29 (transferred to genus *Neomyzomyia*; *in part*, F, Fig. 11); James & Liston, 1911: 106 (F wing).

Female. Head: proboscis length 2.03–2.30 mm (mean 2.20 mm), ratio of length to forefemur length 1.03–1.11 (mean 1.08), maxillary palpus length 1.93–2.20 mm (mean 2.06 mm), ratio of length to proboscis length 0.90–0.99 (mean 0.94), ratio of length to forefemur 0.98–1.02 (mean 1.01), ratio of length of palpomeres 3/4 1.53–1.69 (mean 1.59), 3/5 2.08–2.55 (mean 2.28), 4/5 1.31–1.60 (mean 1.44), 4–5/3 1.00–1.16 (mean 1.07), palpomeres 2–4 with conspicuous apical bands of silvery white scales, pale scales of palpomere 5 white to pale cream-coloured, length of band of palpomere 5 0.91–1.83 (mean 1.21) length of basal dark band of palpomere 5, ratio of length of apical pale band of palpomere 4 to length of basal dark band of palpomere 5 0.27–0.78 (mean 0.56). Thorax: pleural setae as follow: 2–4 upper proepisternal, 0–4 prespiracular, 3–6 prealar, 3–4 upper mesokatepisternal, 2–4 lower mesokatepisternal, 3–6 upper mesepimeral. Wing: length 2.78–3.67 mm (mean 3.33 mm), pale scales on all veins light cream-coloured, spots on costa, subcosta and R_1 more obviously cream-coloured, those on vein R and posterior veins lighter, not strongly contrasting with others; PHP, HP and PSP spots of costa prominent or reduced but usually present, SP spot always present and prominent, ASP spot always absent, PP spot 0.63–1.36 (mean 1.00) length of SCP spot, AD spot 2.00–3.20 (mean 2.43) length of PP spot; PSD spot of vein R extending basally from level 2 to level 5 on one or both wings, PSD spot of vein R with 1–4 pale interruptions, sum of pale interruptions on PSD-PD spots of vein R 4–9 for each wing, ratio of length of cell R_2 to vein R_{2+3} 1.53–1.98 (mean 1.85), ratio of length of cell R_2 to cell M_{1+2} 1.18–1.28 (mean 1.22). Legs: foretarsomeres 2, 3 and 4 with broad basal and apical bands of white scales on dorsal, lateral and anterior surfaces, tan to light brown-scaled along ventral surface, apical and basal pale bands indistinct or absent on ventral surface, foretarsomere 2 sometimes mostly

pale-scaled on dorsal, lateral and anterior surfaces, with 2 narrow dark middle bands, foretarsomere 5 mostly dark-scaled with pale scales at apex, occasionally entirely pale-scaled, dark scales on foretarsomeres 4, 5 sometimes tan to light brown; midtarsomeres 2–4 with apical pale band on dorsal, lateral and anterior surfaces, ventral surface entirely dark-scaled, midtarsomere 5 dark-scaled; hindtarsomeres 2, 3 with short apical pale band, hindtarsomere 2 with 1–3 pale patches on middle dark region, hindtarsomere 4 with basal and apical pale bands, hindtarsomere 5 dark with pale scales at apex. Abdomen: tergum VI without scales, tergum VII with few cream-coloured to brownish scales posteriorly, tergum VIII with cream-coloured to golden scales covering apical 0.5–0.8; sternum VI usually without scales, sometimes with 1–3 dark posteromedial scales, sternum VII with posteromedial patch of dark scales, sternum VIII with small basolateral patch of pale scales.

Male. Essentially as in female except for sexual characters. Wing generally paler, with reduced scaling, pale spots usually longer than in female. Palpomere 2 with dorsal patch of pale scales at middle extending laterally, apex bare; palpomere 3 with long dorsal patch of pale scales at middle, extending laterally, apex with broad band of pale scales covering dorsal, lateral and ventral surfaces, and dark scales at apex of ventrolateral surface, palpomeres 4 and 5 mostly pale-scaled with narrow basal band of dark scales, both with longitudinal line of dark scales on ventral surface not reaching apex of palpomere 5. Abdomen: sternum VI and VII with posteromedial patches of dark scales, sternum VIII covered with pale cream-coloured to yellowish scales.

Pupa (Fig. 5, Table 9). Position and development of setae as figured; range and modal number of branches in Table 9. Measurements and counts from 18 to 20 specimens. In general similar to *An. latens* except for the following characters. Sterna II–VII with narrow dark bands near anterior margin. Cephalothorax: seta 5-CT with 3–6 branches, 6-CT single or double, 7,8,12-CT usually double (7, 2–4; 8,12, 1–3), 9-CT single or double. Seta 9-II–VIII light to medium brown, slightly darker at base. Abdomen: seta 6-I usually double (1,2), 7-I with 2–4 branches; 6-II single or double, 7-II with 3–5 branches, 8,10,11-II absent, 9-II length 0.01–0.014 mm (mean 0.01 mm); 1-III with 4–6 branches, 9-III length 0.02–0.024 mm (mean 0.02 mm); 1-IV with 2–6 branches, 5-IV with 4–6 branches, 6-IV single, 9-IV length 0.02–0.05 mm (mean 0.03 mm), ratios of length of seta 9-IV/9-III 1.33–2.44 (mean 1.82) and 9-IV/9-V 0.23–0.46 (mean 0.33); 1-V usually double (1–3), 5-V with 4–7 branches, 6-V single, 9-V length 0.07–0.13 mm (mean 0.11 mm), with or without minute denticles; 1-VI usually single (1–3), 6-VI single or double, 9-VI length 0.10–0.13 mm (mean 0.11 mm), with or without minute denticles; 1-VII single or double, 5-VII with 3–8 branches, 9-VII length 0.11–0.14 mm (mean 0.12 mm), with or without minute denticles; 9-VIII with 10–20 branches. Paddle: filamentous spicules on outer apical margin and inner margin usually present and prominent, rarely absent on inner margin or from both outer and inner margins; seta 2-Pa single or double; toothed margin index 0.78–0.90 (mean 0.85), paddle teeth well developed, tapering to acute apex, teeth widely spaced.

Larva (Fig. 5, Table 10). Position and development of branches as figured; range and modal number of branches in Table 10. Measurements and counts from 20 to 32 specimens unless otherwise indicated. In general similar to *An. latens* except for the following characters. Head: integument without pattern of dark spots; length 0.68–0.72 mm (mean 0.70 mm) (n 9), width 0.70–0.77 mm (mean 0.74 mm) (n 9), antenna length 0.28–0.32 mm (mean 0.30 mm), ratio of distance between base of seta 1-A to antenna length 0.31–0.44 (mean 0.38); seta 2-C single, simple; 3-C length 0.07–0.10 mm (mean 0.09 mm); distance between bases of 2-C and 3-C 0.03–0.04 mm (mean 0.04 mm), 4-C single, length 0.07–0.10 mm (mean 0.08 mm), extending nearly or beyond base of 2-C, distance between insertions of 2-C and 4-C 0.08–0.11 mm (mean 0.09 mm), ratio of length of 4-C to distance between insertions of 2-C and 4-C 0.80–1.09 (mean 0.93) (n 19); distance between bases of 3-C and 4-C 0.07–0.10 mm (mean 0.08); 5-C as long or slightly longer than antenna, not reaching anterior margin of head, with 5–10 branches, 6-C with 7–13 branches, 7-C with 7–11 branches. Thorax: tubercles of all large setae light brown to yellowish; seta 1-P with 11–17 branches, tubercle of seta 1-P partially joined to tubercle of 2,3-P by an anterior basal bridge, each tubercle with strong pointed apical tooth arising from posterodorsal side and projecting forward over bases of 1,2-P, 14-P with 4–7 branches; 4-M single to triple, 6-M triple, 14-M with 5–9 branches; 3-T moderately developed, palmate, with 3–6 leaflets arising from long slender stem. Abdomen: seta 1-I moderately developed, with 3–6 leaflets, 2-I single to triple, 9-I with 3,4 branches; 1-II moderately developed with 6–10 leaflets; 2-IV with 3,4 branches, 3-IV double or triple, 13-IV triple to 4-branched, ratio of length to length of 10-IV 0.53–0.87 (mean 0.67); 2-V with 4,5 branches; 1-VII with 9–13 leaflets, leaflets without apicolateral serrations; pecten with 12–15 teeth, 4–7 long teeth alternating with 7–10 short teeth.

Material examined. HOLOTYPE: adult pinned female with the head detached and glued on the same round label with the adult. There are four labels with the specimen: a handwritten label ('Kârwâr, Bombay'); a round, handwritten label ('49, *An. Karwar*, 26.4.02'); a small round type label with a red circle; and a rectangular label ('*Anopheles elegans*, Holotype, P.F. Mattingly, 1.ii.1955'). Total material of *An. elegans* examined: 1155 specimens comprising 176F, 203M, 362Le, 369Pe, 45L, derived from three separate collections from natural habitats (2 adult, 1 immature) and a series from colony.

Distribution. *Anopheles elegans* is apparently restricted to peninsular India, where it occurs in hill forests of Karnataka and Tamil Nadu, and was known as the Western Ghat form of *An. dirus* (Bhat, 1988).

Bionomics. Immatures of *An. elegans* were found in a jungle spring in association with *An. culiciformis* Cogill and *An. fluviatilis* James in April (Cogill, 1903), and in elephant footprints in association with *An. mirans*. They

were also found in streams, muddy pools, spring pools and tree holes in the Nilgiris District, India (Tewari *et al.*, 1987). Larval habitats were inside the forest and also in betel nut gardens (Bhat, 1988). Adults were collected in human-bait collections, resting on vegetation and in cattle sheds in the Kyasanur area of Shimoga District, Karnataka (Tewari *et al.*, 1987).

Systematics and nomenclature. James (in Theobald, 1903) described *An. elegans* from an adult female that Cogill had collected in India the year before (Kârwâr, Karnataka). James placed the species in the genus *Myzomyia* and designated the single female he sent to Theobald as 'the type' of *An. elegans* and reported (p. 54) that it was deposited in the 'British Museum (Nat. Hist.), presented by the describer.' James also wrote that neither the male nor the immature stages were known. In October of the same year, Cogill (1903) placed *Myzomyia elegans* in the genus *Anopheles* and mentioned that two larvae of '*A. leucosphyrus*, Dönitz, var. *elegans*, Theobald' were collected in India. The larvae were collected in a jungle stream in April and no additional larvae were found again until the rains ended. Cogill also described the larva, pupa and adult male, and redescribed the female. It is significant that the wing illustration in Cogill's paper differs from that in Theobald (1903), which is *An. elegans*. Cogill's illustration probably belongs to one of the specimens collected in September–October James & Liston (1904) considered *An. elegans* to be a variety of *An. leucosphyrus*, included it in their Group II with *An. punctulatus*, and confirmed that the only specimens available were collected by Cogill in Kârwâr, India. Two wing illustrations appear in James & Liston (1904). The wing illustrated in Plate XI(I) is similar to what Cogill (1903) published and is thus distinct from the wing in Theobald (1903). However, the wing depicted on p. 83 of James & Liston (1904) is identical to that shown on p. 53 in Theobald (1903). In 1910, Theobald described a new genus, *Neomyzomyia*, and transferred *An. elegans* to it. He also redescribed *An. elegans* (because he considered James's description to be inadequate) giving photographs of wings (as Figs 11 and 12) that differ from that of Theobald (1903), especially in having three pale interruptions on the PD spot of vein R1, and in having the PSD spot of vein R with at least one pale interruption (the other pale interruption is not clear in the picture). Moreover, the wings in Figs 11 and 12 are not the same: the latter belongs to a specimen of what appears to be *An. baimaii* (as described herein) collected by R. White on the Andaman Islands. James & Liston (1911) agreed with Theobald (1910) in considering *An. elegans* to be a species of *Neomyzomyia* and reprinted James's original description and wing drawing with some additions from Theobald (1910). Christophers (1933) treated *An. elegans* as a junior synonym of *An. leucosphyrus* and confirmed that the type was in the BMNH. Reid (1949), however, found no specimen labelled as type among Cogill's specimens deposited in the BMNH but confirmed that one female had the same collection data reported by James in Theobald (1903). The other seven specimens in the BMNH were collected by Cogill in September–October 1902. In the NMNH, there




Fig. 5. *Anopheles elegans*. Pupa (female): CT, cephalothorax; Pa, paddle; I-IX, abdominal segments of pupa (dorsal on left, ventral on right), numbers on the left side denote seta 9 of designated segments. Larva: C, head; P, prothorax; M, mesothorax; T, metathorax; PP, pecten plate; abdominal segments I-VI left side dorsal, right side ventral; VII-X lateral (left side) view. Circle shows basal details of setae 1-3-P. Millimetre scale bars.

Table 9. Number of branches for pupal setae of *Anopheles elegans*: range (mode). n.c. = not counted.

Seta no.	Cephalothorax CT	Abdominal segments									Paddle P
		I	II	III	IV	V	VI	VII	VIII	IX	
0	-	-	1	1	1	1	1	1	1	-	-
1	1-3 (2)	n.c.	>30	4-6 (5)	2-6 (3)	1-3 (2)	1-3 (1)	1,2 (1)	-	1	1
2	1-3 (2)	6-8 (7)	4-6 (5)	4-7 (6)	2-4 (3)	2-4 (3)	2,3 (3)	2,3 (3)	-	-	1,2 (1)
3	2-3 (3)	1	1,2 (2)	2-4 (2)	5-7 (6)	1-3 (2)	1,2 (2)	2,3 (3)	-	-	-
4	1-3 (2)	4-6 (4)	4-6 (5)	2-6 (4)	3-5 (4)	2-5 (4)	1,2 (2)	1,2 (2)	1-3 (2)	-	-
5	3-6 (5)	2,3 (2)	2-6 (3)	6-11 (6)	4-6 (5)	4-7 (5)	3-7 (4)	3-8 (5)	-	-	-
6	1,2 (2)	1,2 (2)	1,2 (2)	1,2 (2)	1	1	1,2 (1)	1,2 (2)	-	-	-
7	2-4 (2)	2-4 (3)	3-5 (3)	1-4 (3)	2-4 (3)	2-4 (2)	1	1	-	-	-
8	1-3 (2)	-	-	1-4 (3)	1-3 (3)	1-3 (2)	1-3 (2)	1-3 (2)	-	-	-
9	1,2 (1)	1	1	1	1	1	1	1	10-20 (11)	-	-
10	2-4 (3)	-	-	1-3 (2)	1,2 (1)	1,2 (2)	1,2 (2)	1-4 (2)	-	-	-
11	2-4 (3)	-	-	1	1	1	1	1,2 (2)	-	-	-
12	1-3 (2)	-	-	-	-	-	-	-	-	-	-
13	-	-	-	-	-	-	-	-	-	-	-
14	-	-	-	-	1	1	1	1	1	-	-

Table 10. Number of branches for setae of fourth-instar larva of *An. elegans*: range (mode).

Seta no.	Head C	Thorax			Abdominal segments								
		P	M	T	I	II	III	IV	V	VI	VII	VIII	X
0	1	1	-	-	-	1	1	1	1	1	1	1	-
1	1	11-17 (13)	23-33 (24)	1	3-6 (5)	6-10 (7)	17-22 (20)	17-22 (20)	18-21 (19)	16-20 (18)	9-13 (13)	1	1
2	1	9-15 (11)	1-4 (2)	1	1-3 (2)	5-9 (7)	5-8 (6)	3-4 (4)	4-5 (4)	4-6 (4)	5-8 (7)	4-8 (6)	20
3	1	1	1	3-6 (5)	1-2 (1)	1	1	2-3 (3)	1	1	1-3 (2)	3-5 (4)	8
4	1	13-22 (15)	1-3 (2)	2-4 (3)	3-7 (4)	4-9 (4)	2-4 (3)	3-4 (3)	1-3 (3)	1	1	1	9
5	5-10 (8)	n.c.	1	24-30 (28)	2-3 (3)	3-5 (4)	3-5 (4)	3-4 (3)	3-5 (4)	4-6 (4)	3-5 (4)	2-4 (3)	-
6	7-13 (10)	1	3	2-3 (2)	15-20 (15)	17-23	13-16 (15)	2-3 (2)	2-3 (2)	3	3-5 (4)	1-S	3-7 (4)
7	7-11 (9)	14-25 (18)	2-3 (2)	22-28	18	n.c.	3-5 (4)	2-5 (5)	3-4 (4)	2	1-3 (2)	2-S	4-7 (5)
8	1	20-26 (26)	n.c.	n.c.	-	1-2 (1)	1-3 (2)	1-3 (2)	2-5 (2)	2-3 (2)	3-8 (4)	6-S	2-3 (2)
9	1-3 (2)	1	1	1	3-4 (3)	5-8 (6)	6-10 (8)	7-9 (8)	6-8 (7)	6-7 (7)	2-5 (3)	7-S	1
10	1-2 (1)	1	1	1	1	2-4 (2)	1	1	1	2-4 (3)	3-5 (4)	8-S	2-3 (2)
11	n.c.	1	1	1	3	1	2-4 (2)	2-3 (2)	2-3 (2)	2	2	9-S	2-4 (3)
12	1-2 (1)	1-3 (1)	1	2	2-3 (3)	1	2-3 (2)	2-3 (2)	1-3 (2)	1	1-2 (1)	-	-
13	4-5 (4)	3-5 (4)	3-6 (5)	2-3 (3)	3-5 (3)	3-6 (4)	4-7 (4)	3-4 (3)	2-3 (3)	4-7 (5)	2	-	-
14	n.c.	4-7 (4)	5-9 (5)	-	-	-	-	1	1	1	1	1	-
15	5-6 (6)	-	-	-	-	-	-	-	-	-	-	-	-

are also four of Cogill's specimens collected in September–October 1902. Reid (1949) concluded that the description of James & Liston (1904) was based on the specimens collected by Cogill in September–October 1902 and suggested further that James in Theobald (1903) was wrong in designating the specimen collected in April 1902 as the type of *An. elegans*. In a discussion on the validity of the name *An. elegans*, Colless (1956) considered the single female (collected by Cogill, v.02) James sent to Theobald to be '*An. l. balabacensis*', and the other specimens to be *An. elegans*. He concluded this because Theobald did not label the single female as the type and because James & Liston's (1904, 1911) descriptions were probably based on specimens collected in September–October. Colless also considered Theobald's designation of the single female collected in April as the 'Type' was an error and the remaining four females and two males (collected by Cogill, x.02) should be considered cotypes, from which a lectotype should be designated.

A holotype is defined as 'the single specimen upon which a new nominal species-group taxon is based in the original publication' (Article 73.1: ICZN, 1999). Article 73.1.1 also says 'If an author when establishing a new nominal species-group taxon states in the original publication that one specimen, and only one, is the holotype, or "the type", or uses some equivalent expression, that specimen is the holotype fixed by original designation'. Article 73.1.3 states, 'The holotype of a new nominal species-group taxon can only be fixed in the original publication and by the original author.' From these three articles, it is clear that the specimen designated as 'the type' by James in Theobald (1903) is the holotype of *An. elegans*. In addition, James clearly wrote that 'the specimen is imperfect' and was collected in April. Since James only sent a single female and only this female was noted in the original description, that specimen is the holotype. Colless (1956) examined Cogill's specimens of *An. elegans* (i.e. original specimens), as well as the correspondence between James and Theobald, and concluded that James sent Theobald a written description, figure of one wing and one female with the suggestion that this specimen might be kept as a type until better specimens could be sent. Examination of the specimen designated as 'the type' by James revealed that it is imperfect, with the head detached from the body and glued to the same label with the pinned adult female. Mattingly recognized the specimen as the holotype of *An. elegans* and added a type label and an additional label on which he wrote '*Anopheles elegans* James, Holotype, P.F. Mattingly, 1.ii.1955. We examined the 12 specimens collected by Cogill in 1902, both in April and September–October 1902 (BMNH and NMNH) and discovered that they belong to two distinct species of the Leucosphyrus Group. Because the holotype of *An. elegans* is the specimen designated as the type by James in Theobald (1903), the other specimens collected in September and October belong to an undescribed species that has been largely misidentified as *An. elegans* since Cogill (1903). Furthermore, the holotype shows that *An. elegans* belongs to the Dirus Complex of the Leucosphyrus Subgroup, whereas the undescribed species belongs to the Elegans

Subgroup of Colless (1957) and Peyton (1990). Also, *Anopheles dirus* species E (Sawadipanich *et al.*, 1990) is, in fact, *An. elegans*. Because *An. elegans* belongs to the Dirus Complex, the group designated the Elegans Subgroup by Colless and Peyton is now renamed as the Hackeri Group.

Cytogenetic and crossing data confirm that *An. elegans* (as *An. dirus* species E) is a distinct species in the Dirus Complex and is more closely related genetically to *An. baimaii* than to any other member of the complex. Both species, for example, share the same Xa inversion. It seems that *An. elegans* and *An. baimaii* evolved from a common ancestor through allopatric speciation (Sawadipanich *et al.*, 1990).

Adults of *An. elegans* are distinguished, except from *An. scanloni* and *An. baimaii*, by the following combination of characters: (1) proboscis dark-scaled, slightly longer than forefemur, ratio of length to forefemur length 1.03–1.11; (2) PSD spot of vein R rarely extending to level 1, 2 or 3 on at least one wing, frequently extending to level 4 and 5 on one or both wings; (3) PSD spot of vein R usually with 2–4 pale interruptions on at least one wing, rarely with one pale interruption on one wing; (4) ASP spot absent on costa and usually absent on subcosta; (5) hindtarsomere 4 with basal and apical pale bands; (6) hindtarsomere 5 dark-scaled at base; (7) apical pale band on hindtibia entirely white, without dark extension; (8) vein 1A with a long pale spot at level of the PSD spot of vein R on at least one wing, this pale spot sometimes reduced but always the longest pale spot of 1A; (9) sternum VI usually without scales, occasionally with 1–3 posteromedial dark scales.

The fourth-instar larva of *An. elegans* can be recognized by the following combination of characters: (1) seta 5-C as long as or slightly longer than the antenna, not reaching the anterior margin of the head; (2) seta 1-X inserted on the saddle; (3) seta 2-C without aciculae; (4) tubercle of seta 1-P narrowly joined to the tubercle of 2,3-P by an anterior basal bridge; (5) tubercle of setae 1,2-P with strong, pointed apical tooth arising from posterodorsal side and projecting forward over base of 1,2-P; (6) seta 4-C not reaching the base or extending slightly beyond the base of 2-C, ratio of length of 4-C to distance between 2-C and 4-C 0.80–1.09. Recognition of *An. elegans* based on pupal characters is not reliable. A few individuals examined did not have a fringe of filamentous spicules on the inner margin of the paddle, and thus they could have been misidentified as *An. scanloni*. Because *An. elegans* and *An. scanloni* are allopatric, their geographical distributions can facilitate their identification.

Hackeri subgroup (formerly Elegans subgroup)

Anopheles (Cellia) recens Sallum & Peyton, sp.n.

Etymology: the name *recens* is Latin for fresh, young, recent.

Anopheles Sumatra species of Peyton, 1990: 197.

Female. Head: proboscis uniformly dark-scaled or with long ventral patch of pale scales on apical 0.2 of ventral

surface, length 2.28–2.68 mm (mean 2.50 mm), ratio of length to forefemur length 1.16–1.30 (mean 1.21); maxillary palpus length 1.80–2.33 mm (mean 2.10 mm), ratio of length to proboscis length 0.79–0.86 (mean 0.83), ratio of length to forefemur length 0.97–1.05 (mean 1.01), ratio of length of palpomeres 3/4 1.67–2.54 (mean 1.94), 3/5 2.50–3.62 (mean 3.05), 4/5 1.22–2.12 (mean 1.60), 4–5/3 0.70–1.00 (mean 0.85), palpomeres 2–4 with narrow apical pale cream-coloured bands, pale scales on palpomere 5 almost yellowish, apical pale band of palpomere 5 0.00–1.00 (mean 0.5) length of basal dark band of palpomere 5, ratio of length of apical pale band of palpomere 4 to length of basal dark band of palpomere 5 0.00–0.67 (mean 0.21). Thorax: pleural setae as follows: 2–5 upper proepisternal, 0–2 prespiracular, 5–9 prealar, 6,7 upper mesokatepisternal, 3–6 lower mesokatepisternal, 3–8 upper mesepimeral. Wing: length 3.64–4.55 mm (mean 4.05 mm), pale scales on all veins light cream-coloured, spots on subcosta and SCP, PP and AP spots of costa more obviously cream-coloured nearly yellowish, HP spot present, prominent, PSP spot rarely absent, SP spot usually present, prominent, ASP spot absent on costa, PP spot 0.79–3.80 (mean 1.31) length of SCP spot, AD spot 1.05–3.63 (mean 2.58) length of PP spot, PSD spot of vein R extending basally from level 2 to level 4 on both wings, PSD spot of vein R with 1–4 pale spots on one or both wings, SD spot of vein R usually with 1, 2 pale interruptions, rarely with 3 pale interruptions on both wings, sum of pale interruptions on PSD-PD spots of vein R 3–8 for each wing; ratio of length of cell R_2 to vein R_{2+3} 1.36–1.79 (mean 1.58), ratio of length of cell R_2 to cell M_{1+2} 1.27–1.48 (mean 1.36). Legs: foretarsomeres 2,3 with short basal and apical bands of pale scales, foretarsomere 2 with 2–4 pale spots on middle of dorsal surface, foretarsomere 4 with indistinct basal and apical pale bands, usually without basal pale band, foretarsomere 5 with pale scales at apex or entirely dark-scaled, foretarsomeres 2–4 entirely dark-scaled on ventral surface; midtarsomeres 2–4 with small, apical patches of pale scales on dorsal surface, midtarsomeres 2,3 with 1–3 pale spots on middle dark area, dark-scaled ventrally, midtarsomere 5 entirely dark-scaled; hindtarsomeres 2–4 with short apical pale bands, hindtarsomere 2 with 2–4 pale spots on median dark area, hindtarsomere 5 dark-scaled. Abdomen: tergum VI without scales, tergum VII with 2–4 dark scales on posterior margin, tergum VIII with patch of dark scales medially, few golden scales posterolaterally; sternum VI usually without scales, rarely with 2,3 posteromedial scales, sternum VII without scales or with posteromedial patch of dark scales, sternum VIII without scales, rarely with 2,3 anterolateral pale scales.

Male. Essentially as in female except for sexual characters. Wing generally paler with reduced scaling, pale spots usually longer than in female. Proboscis dark-scaled without ventral patch of pale scales; palpomere 2 with dorsal patch of pale scales at middle, apex of palpomere 2 bare, palpomere 3 with long dorsal patch of pale scales at middle extending laterally, sometimes reaching ventral surface, apex of palpomere 3 variable, with broad band of pale

scales covering dorsal and lateral surfaces, with dorsal patch of pale scales, or entirely dark-scaled, palpomere 4 variable, pale-scaled with basal band of dark scales, less frequently with patch of dark scales on ventral and dorsal surfaces, palpomere 5 variable, pale-scaled with basal dark, or with basal dark band with extension along ventral surface not reaching apex of segment, or mostly dark-scaled with lateral patch of pale scales. Abdomen: sternum VIII covered with pale cream-coloured to whitish scales.

Pupa (Fig. 6, Table 11). Position and development of setae as figured; range and modal number of branches in Table 11. Measurements and counts from 7 to 10 specimens. In general similar to *An. latens* except for the following characters. Sterna II–VI with narrow dark band near anterior margin. Cephalothorax: seta 5-CT with 3–5 branches, 6-CT single or double, 7–12-CT usually double (see Table 11 for ranges). Seta 9-IV–VIII light to medium brown, slightly darker at base. Abdomen: seta 6-I usually single (1,2), 7-I with 2–4 branches; 1-II dendritic, with 8–18 narrow branches, 6-II single or double, 7-II double or triple, 8,10,11-II absent, 9-II length 0.01–0.02 mm (mean 0.02 mm); 1-III with 3–5 branches, 5-III with 5–8 branches, 6-III usually single (1,2), 9-III length 0.02–0.03 mm (mean 0.02 mm); 1-IV with 2–4 branches, 5-IV with 4,5 branches, 6-IV single, 9-IV long, always simple, length 0.05–0.08 mm (mean 0.06 mm), ratios of length of 9-IV/9-III 1.92–3.26 (mean 2.27) and 9-IV/9-V 0.33–0.76 (mean 0.44); 1-V usually double (1,2), 5-V with 4–6 branches, 6-V single, 9-V long, always simple, length 0.11–0.14 mm (mean 0.13 mm); 1-VI single or double, 5-VI with 3–5 branches, 6-VI single, 9-VI always simple, length 0.12–0.15 mm (mean 0.13 mm); 1-VII single, 5-VII with 4–6 branches, 6-VII single, 9-VII always simple, length 0.12–1.15 mm (mean 0.14 mm); 9-VIII with 6–13 branches. Paddle: seta 2-Pa single; toothed margin index 0.84–0.93 (mean 0.89), paddle teeth well developed, tapering to apex, ending in acute apex, widely spaced.

Larva (Fig. 6, Table 12). Position and development of setae as figured; range and modal number of branches in Table 12. Measurements and counts from 8 to 16 specimens unless otherwise indicated. In general similar to *An. latens* except for the following characters. Head: integument light brown to yellowish with dark spots on dorsal apotome and labiogula as follow: one dark spot centrally on dorsal apotome, posterior to setae 5,6-C; a more prominent, single dark spot posterior to area between seta 8-C and 2 small dark spots laterally; 2 lateral dark spots at dorsal apotome, posterior to seta 7-C; 1 dark spot at ventral lateral area near insertion of seta 12-C; 2 dark spots placed laterally at the level of posterior tentorial pit, and a single dark spot posterior on hypostomal ecdysial line; head length 0.75–0.81 mm (mean 0.79 mm) (n 5), width 0.73–0.83 mm (mean 0.76 mm) (n 5); antenna length 0.33–0.40 mm (mean 0.37 mm), ratio of distance between base and seta 1-A to antenna length 0.32–0.43 (mean 0.38); seta 2-C usually rarely with few, sparse, minute spicules; 3-C length 0.11–0.15 mm (mean 0.13 mm); distance between bases of 2-C and 3-C 0.04–0.06 mm




Fig. 6. *Anopheles recens* sp.n. Pupa (female): CT, cephalothorax; Pa, paddle; I-IX, abdominal segments of pupa (dorsal on left, ventral on right), numbers on the left side denote seta 9 of designated segments. Larva: C, head; P, prothorax; M, mesothorax; T, metathorax; PP, pecten plate; abdominal segments I-VI left side dorsal, right side ventral; VII-X lateral (left side) view. Circle shows basal details of setae 1-3-P. Millimetre scale bars.

Table 11. Number of branches for pupal setae of *Anopheles recens*: range (mode).

Seta no.	Cephalothorax CT	Abdominal segments									Paddle P
		I	II	III	IV	V	VI	VII	VIII	IX	
0	—	—	1	1	1	1	1	1	1	—	—
1	2	9–16 (12)	8–18 (12)	3–5 (4)	2–4 (3)	1–2 (2)	1–2 (1)	1	—	1	1
2	2	3–5 (4)	2–4 (4)	2–5 (3)	2–3 (2)	1–2 (2)	1–2 (2)	2–3 (2)	—	—	1
3	2	1	1–2 (2)	1–3 (2)	2–5 (4)	1–2 (2)	1–2 (2)	2–3 (3)	—	—	—
4	1–2 (2)	2–4 (3)	2–4 (2)	2–4 (3)	2–4 (3)	2–4 (3)	1–2 (2)	1–2 (2)	1–2 (1)	—	—
5	3–5 (4)	2–5 (3)	2–4 (2)	5–8 (6)	4–5 (4)	4–6 (4)	3–5 (4)	4–6 (4)	—	—	—
6	1–2 (1)	1–2 (1)	1–2 (1)	1–2 (1)	1	1	1	1	—	—	—
7	2–4 (2)	2–4 (3)	2–3 (3)	1–3 (2)	2–3 (2)	2–3 (2)	1–2 (1)	1	—	—	—
8	1–3 (2)	—	—	2–3 (3)	1–3 (2)	1–3 (2)	2–3 (2)	1–2 (2)	—	—	—
9	1–2 (2)	1	1	1	1	1	1	1	6–13 (7)	—	—
10	2–3 (2)	—	—	2–3 (2)	1–2 (1)	1–2 (1)	1–2 (2)	1–2 (2)	—	—	—
11	1–4 (2)	—	—	1	1	1	1	1–2 (1)	—	—	—
12	2–3 (2)	—	—	—	—	—	—	—	—	—	—
13	—	—	—	—	—	—	—	—	—	—	—
14	—	—	—	—	1	1	1	1	1	—	—

Table 12. Number of branches for setae of fourth-instar larva of *An. recens*: range (mode).

Seta no.	Head C	Thorax			Abdominal segments								
		P	M	T	I	II	III	IV	V	VI	VII	VIII	X
0	1	1	—	—	—	1	1	1	1	1	1	1	—
1	1	14–20 (16)	23–30 (26)	1	3–4 (4)	5–8 (5)	16–23 (18)	13–22 (16)	15–21 (18)	14–20 (16)	8–12 (8)	1	1–2 (1)
2	1	11–14 (11)	1–3 (1)	1	1	4–8 (5)	4–6 (5)	3	3–5 (3)	3–5 (4)	4–6 (5)	3–5 (4)	11–16 (14)
3	1	1	1	3–6 (5)	1	1	1	1–2 (1)	1	1–2 (1)	1–3 (2)	3–5 (4)	5–7 (6)
4	1	10–15 (14)	1–2 (1)	2	3–5 (3)	3–4 (4)	1–3 (2)	1–3 (2)	1–3 (3)	1	1	1	9
5	9–12 (12)	16–28 (25)	1	26–34	3	3–5 (4)	3–4 (3)	3	2–3 (3)	3–5 (3)	2–4 (3)	2–5 (3)	—
6	10–13 (12)	1	3–4 (4)	2–3 (3)	12–17 (14)	12–18 (14)	10–16 (12)	2–3 (2)	1–2 (2)	1–3 (2)	2–4 (2)	1–5 (4)	4–6 (4)
7	11–15 (13)	16–23 (21)	2–4 (2)	23–27 (25)	9–14 (11)	10–15 (13)	2–4 (3)	1–5 (3)	2–4 (3)	2	2–3 (3)	2–5 (4)	4–8 (5)
8	1–2 (1)	16–30 (17)	16–23 (16)	22–30 (30)	—	1–2 (2)	1–2 (1)	1–2 (2)	2	1–2 (2)	2–4 (3)	6–8 (3)	2–3 (2)
9	2–4 (2)	1	1	1	3–4 (3)	4–8 (7)	4–8 (7)	5–8 (5)	4–7 (4)	4–9 (5)	3–4 (3)	7–9 (3)	1–2 (2)
10	1	1	1	1	1	1–3 (2)	1	1	1	2–3 (2)	4–5 (4)	8–9 (2)	1–2 (2)
11	16–24 (17)	1	1	1	3	1–2 (1)	2	2	2–3 (2)	1–3 (1)	1–2 (1)	1–3 (1)	—
12	1–2 (2)	1	1	2–3 (3)	3	1	1–2 (2)	1–2 (2)	1–2 (1)	1	1	—	—
13	3–5 (4)	3–5 (4)	3–5 (5)	1–3 (2)	3–6 (4)	2–4 (3)	2–4 (3)	1–2 (2)	2	4–8 (4)	2	—	—
14	3–6 (4)	5–9 (8)	4–7 (5)	—	—	—	—	1	1	1	1	1	—
15	4–6 (4)	—	—	—	—	—	—	—	—	—	—	—	—

(mean 0.05 mm); 4-C length 0.12–0.17 mm (mean 0.15 mm), extending beyond base of 2-C, distance between insertions of 2-C and 4-C 0.08–0.11 mm (mean 0.09 mm), ratio of length of 4-C to distance between insertions of 2-C and 4-C 1.44–2.0 (mean 1.68), distance between bases of 3-C and 4-C 0.05–0.08 mm (mean 0.07 mm), 5-C shorter than antenna, not reaching anterior margin of head, with 9–12 branches, 6-C with 10–13 branches, 7-C with 11–15 branches. Thorax: tubercles of all large setae light to medium brown; seta 1-P with 14–20 branches, tubercle of seta 1-P usually joined basally to tubercle of 2,3-P, rarely separate, both tubercles with strong, pointed or somewhat rounded posterodorsal tooth projecting forward over base of 1,2-P, 14-P with 5–9 branches; 4-M single or double, 6-M usually 4-branched (3,4), 14-M with 4–7 branches; 3-T somewhat palmate, with 3–6 leaflets. Abdomen: seta 1-I not palmate, with 3,4 slender branches, 2-I always single, 9-I with 3,4 branches; 1-II moderately developed, more pigmented than 1-I, somewhat palmate, with 5–8 leaflets; 2-IV always triple, 3,13-IV single or double, ratio of length of 13-IV to 10-IV 0.91–1.24 (mean 1.11) (n 7); 1-VII with 8–12 leaflets, leaflets with minute apicolateral serrations and slender apical filament; 1-X single or double, inserted on saddle; pecten teeth 14–17 teeth, 5–8 long teeth alternating with 6–11 short teeth.

Material examined. HOLOTYPE: adult female with associated larval and pupal exuviae on microscope slide, acc. no. IN704-10, collected by U. Malaya team, 5 April 1978, deposited in the NMNH. Type locality: INDONESIA, Sumatra Island, Gunong Dempo (4°2'S 103°9'E). PARATYPES: same data as holotype, 8F, 11M, 6Le, 6Pe, 7L; acc. no. IN705, 5 April 1978, 2F, 1M. Other material of *An. recens* examined: 44 specimens comprising 11F, 12M, 7Le, 7Pe, 7L derived from 2 separate collections from natural habitats (2 immatures).

Distribution. *Anopheles recens* is known only from Sumatra Island, Indonesia.

Bionomics. Nothing is known about the medical importance of *An. recens* or the involvement of this species in the transmission of parasites to humans and other animals. Immatures of *An. recens* were taken from temporary ground pools in cloud forests. The water was fresh, stagnant or slow moving, clear with or without emergent vegetation, and in partial shade or full sun. The larval habitats were located at elevations of 1550 m above sea level.

Systematics. *Anopheles recens* (as *An. leucosphyrus* Sumatra species) was included in the Hackeri Group (as *Elegans* Group) because the proboscis is noticeably longer than the forefemur and the palpus is shorter than the proboscis (Peyton, 1990). Adults can be distinguished by the following characters: (1) proboscis longer than forefemur, ratio of proboscis length to forefemur length 1.16–1.30 (mean = 1.20); (2) proboscis of female entirely dark-scaled or with ventral patch of pale scales on apical 0.2; (3) apical pale band of hindtibia entirely white-scaled or with a patch of

dark scales on anterior surface; (4) palpomere 5 entirely dark-scaled or with apical pale band, ratio of length of apical pale band to length of basal dark band 0.00–1.00 (mean = 0.50); (5) PSD spot of vein R extending from level 1 to level 4; (6) PSD spot of vein R with 1–4 pale interruptions; (7) sum of pale interruptions on PSD-PD spots of vein R 3–8 for each wing; (8) SCP, PP and AP spots of costa cream-coloured, almost yellowish, strongly contrasting with other pale spots; (9) tergum VIII with a patch of dark scales medially and a few golden scales posterolaterally; (10) hindtarsomeres without pale scales at base; (11) ASP spot absent from costa; and (12) pale spots on costa small.

The fourth-instar larva of *An. recens* can be recognized by the following combination of characters: (1) seta 5-C conspicuously longer than antenna; (2) basal sclerotized tubercles of setae 1,2-P with prominent tooth or spine projecting from posterodorsal margin forward over the base of each seta; (3) seta 3-C long, extending beyond anterior margin of head; (4) seta 4-C always extending noticeably beyond base of 2-C; (5) seta 1-II with 5–8 very narrow leaflets, leaflets translucent or very lightly pigmented, stem not inflated; (6) seta 13-IV single or double, about length of 10-IV, ratio of length of 13-IV to 10-IV 0.91–1.24 (mean = 1.11); (7) seta 13-V double; (8) seta 2-VII with 4–6 branches; (9) seta 9-VII with 3,4 branches.

Anopheles (Cellia) mirans Sallum & Peyton, sp.n.

Etymology. The name *mirans* is derived from Latin 'mirari' meaning to wonder at, look at.

Anopheles leucosphyrus var. *elegans* of Cogill, 1903: 330 (larval bionomics, in part, F*, M, P, L*); James & Liston, 1904: 82 (F*, L*, in part, Plate XII, transferred to genus *Anopheles*); Reid, 1949: 46 (taxonomic notes, in part, distribution).

Anopheles balabacensis of Reid, 1970: 56 (in part, west coast of India, systematics).

Anopheles elegans of Colless, 1956: 75 (F*, M, P*, E, taxonomic notes, identification key, distribution); Choudhury *et al.*, 1963b: 243 (vector competence); Choudhury *et al.*, 1963a: 237 (vector competence); Reid, 1968: 303 (F*, taxonomy, bionomics notes, identification key); Nelson *et al.*, 1971: 46 (vector competence); Mendis *et al.*, 1984: 318 (L*, P*, bionomics); Peyton, 1990: 197 (taxonomy).

Neomyzomyia elegans of James & Liston, 1911: 105 (F*, L*, Plate XV).

Neomyzomyia leucosphyra of Swellengrebel, 1916: 82 (in part, taxonomy, distribution).

Female. Head: proboscis uniformly dark-scaled, length 2.17–2.86 mm (mean 2.59 mm), ratio of length to forefemur length 1.19–1.31 (mean 1.25), maxillary palpus length 1.85–2.45 mm (mean 2.22 mm), ratio of length to proboscis length 0.84–0.93 (mean 0.87), ratio of length to forefemur

length 1.02–1.15 (mean 1.10), ratio of length of palpomeres 3/4 1.35–1.77 (mean 1.55), 3/5 1.93–2.54 (mean 2.21), 4/5 1.27–1.73 (mean 1.43), 4–5/3 1.04–1.18 (mean 1.10), palpomeres 2–4 with narrow apical silvery white bands, bands similar in size and colour, pale scales of palpomere 5 white, similar to scales on palpomeres 2–4, length of apical pale band of palpomere 5 0.83–2.67 (mean 1.40) length of basal dark band of palpomere 5, ratio of length of apical pale band of palpomere 4 to length of basal dark band of palpomere 5 0.33–1.00 (mean 0.6). Thorax: pleural setae as follows: 2–5 upper proepisternal, 0–1 prespiracular, 4–8 prealar, 3,4 upper mesokatepisternal, 2,3 lower mesokatepisternal, 2–6 upper mesepimeral. Wing: length 2.88–4.23 mm (mean 3.64 mm \pm 0.34), pale scales of PHP, HP, PSP, SP and SCP spots white on costa, SCP spot pale cream-coloured to yellowish on subcosta, pale scales on remaining veins pale cream-coloured to yellowish, none strongly contrasting with others, PHP spot of costa usually prominent, sometimes reduced, HP, PSP and SP spots always present, prominent, ASP spot absent, PP spot 0.46–1.89 (mean 1.00) length of SCP spot, AD spot 1.94–8.00 (mean 3.27) length of PP spot, PSD spot of vein R extending basally from level 1 to level 4, often to level 1 and level 2 on one or both wings, PSD spot of vein R without pale interruptions, SD spot of vein R with 1 pale interruption, sum of pale interruptions on PSD-PD spots of vein R always 2 on both wings, ratio of length of cell R_2 to vein R_{2+3} 1.29–1.80 (mean 1.52), ratio of length of cell R_2 to cell M_{1+2} 1.13–1.29 (mean 1.22). Legs: femora, tibiae and tarsomeres dark-scaled and speckled with pale spots; foretarsomere 2 with conspicuous basal and apical pale bands, middle dark area with 1–3 pale spots or entirely dark, foretarsomere 3 with basal and apical pale bands, foretarsomere 4 with basal and apical pale bands, occasionally apical pale band absent, foretarsomeres 5 dark at base, apex pale-scaled, basal and apical pale bands on foretarsomeres 2–5 evident on ventral surface; midtarsomeres 2–4 with apical pale spot on dorsal surface, pale spots not noticeable on ventral surface, midtarsomere 5 pale-scaled at apex; hindtarsomeres 2,3 with minute basal pale spot or dark-scaled at base, always with apical pale band, hindtarsomere 4 with basal and apical pale bands, hindtarsomere 5 with basal pale band, rarely entirely dark at base, apex always pale-scaled. Abdomen: tergum VI usually without scales, rarely with few dark scales posteriorly, tergum VII usually without scales, occasionally with few apicolateral scales, tergum VIII covered with golden scales posteromedially, dark scales basally; sternum VI with posteromedial patch of dark scales, sternum VII with posteromedial patch of dark scales, sternum VIII with few, occasionally numerous lateral pale cream-coloured scales.

Male. Essentially as in female except for sexual characters. Palpomere 2 with dorsal patch of pale scales at middle, apex of palpomere 2 bare, palpomere 3 with long dorsal patch of pale scales at middle, extending to lateral surface, apex of palpomere 3 with a broad band of pale scales, covering dorsal, lateral and ventral surfaces except for a small ventrolateral patch of dark scales at apex, palpomere

4 mainly pale-scaled with basal dark band extending along ventral surface, palpomere 5 pale-scaled with basal dark band extending along ventral surface and reaching apex. Abdomen: sternum VIII covered with pale cream-coloured to golden scales.

Pupa (Fig. 7, Table 13). Position and development of setae as figured; range and modal number of branches in Table 13. Measurements and counts from 20 to 40 specimens. Sterna II–VI with narrow, dark band near anterior margin. Cephalothorax: seta 4-CT usually double (2–4), 5-CT with 3–7 branches, 6-CT single to triple, 7-CT usually triple (2–4), 8-CT single or double, 10-CT single to triple, 11-CT single to 5-branched. Seta 9-II lightly pigmented, 9-III–VII medium to dark brown from base to apex, 9-VIII light brown, darker at base. Abdomen: seta 6-I single or double, 7-I with 2–5 branches; 6-II single or double, 7-II with 2–5 branches, 9-II length 0.01–0.02 mm (mean 0.01 mm), 10-II single to triple, 11-II absent or present; 1-III with 3–8 branches, 5-III with 5–9 branches, 9-III length 0.01–0.02 mm (mean 0.02 mm); 1-IV with 2–6 branches, 5-IV with 4–7 branches, 6-IV single or double, 9-IV short, either with or without minute denticles, length 0.03–0.06 mm (mean 0.04 mm), ratios of length of 9-IV/9-III 1.41–3.23 (mean 2.32) and 9-IV/9-V 0.29–0.52 (mean 0.41); 1-V usually double (1–3), 5-V with 3–7 branches, 6-V single, 9-V long, either with or without minute denticles, length 0.08–0.12 mm (mean 0.11 mm); 1-VI single or double, 5-VI with 4–7 branches, 6-VI single, 9-VI length 0.07–0.14 mm (mean 0.11 mm); 1-VII single or double, 6-VII usually single, 9-VII length 0.08–0.12 mm (mean 0.11 mm); 9-VIII with 8–13 branches. Paddle: seta 2-Pa with 1,2 branches; toothed margin index 0.78–0.94 (mean 0.85), paddle teeth tapering to a pointed, sharply acute apex.

Larva (Fig. 7, Table 14). Position and development of branches as figured; range and modal number of branches in Table 14. Measurements and counts from 22 to 44 specimens unless otherwise indicated. Head: integument light brown to yellowish with dark spots on dorsal apotome and labiogula as follows: 1 small dark spot at posterior end of dorsal apotome and 2 always slightly anterior; one dark spot posterior to area between seta 8-C and 2 small dark spots laterally; 2 lateral dark spots at dorsal apotome posterior to seta 7-C; 1 dark spot on lateralia in area of insertion of seta 12-C; 2 dark spots laterally at level of posterior tentorial pit; one dark spot placed posteriorly on hypostomal ecdysial line; head length 0.66–0.74 mm (mean 0.72 mm), width 0.70–0.80 mm (mean 0.76 mm), antenna length 0.29–0.33 mm (mean 0.31 mm), ratio of distance between base of seta 1-A to antenna length 0.32–0.50 (mean 0.39); seta 2-C single, occasionally with minute sparse spicules on apical 0.5; 3-C length 0.07–0.10 mm (mean 0.08 mm), distance between bases of 2-C and 3-C 0.03–0.06 mm (mean 0.04 mm), 4-C single to triple, length 0.10–0.16 mm (mean 0.12 mm), extending beyond base of 2-C, distance between basal insertions of 2-C and 4-C 0.07–0.12 mm (mean 0.09 mm), ratio of length of 4-C to distance between insertions of 2-C and 4-C 1.05–2.12 (mean 1.45), distance between bases of 3-C and 4-C




Fig. 7. *Anopheles mirans* sp.n. Pupa (male): CT, cephalothorax; Pa, paddle; I-IX, abdominal segments of pupa (dorsal on left, ventral on right), numbers on the left side denote seta 9 of designated segments. Larva: C, head; P, prothorax; M, mesothorax; T, metathorax; abdominal segments I-VI left side dorsal, right side ventral; VII-X lateral (left side) view. Circle shows basal details of setae 1-3-P. Millimetre scale bars.

Table 13. Number of branches for pupal setae of *Anopheles mirans*: range (mode). n.c. = not counted.

Seta no.	Cephalothorax CT	Abdominal segments									Paddle P
		I	II	III	IV	V	VI	VII	VIII	IX	
0	–	–	1	1	1	1	1	1	1	–	–
1	2	n.c.	11–25	3–8 (5)	2–6 (3)	1–3 (2)	1,2 (1)	1,2 (1)	–	1	1
2	2,3 (2)	3–10 (7)	5–7 (6)	3–8 (6)	2–4 (3)	2–4 (2)	1,2 (1)	2–4 (3)	–	–	1,2 (2)
3	2,3 (2)	1–3 (2)	2–5 (3)	4–10 (5)	3–7 (5)	2,3 (2)	1–4 (2)	2–4 (3)	–	–	–
4	2–4 (2)	2–8 (4)	2–5 (3)	2–4 (3)	2–8 (4)	2–5 (3)	1–3 (2)	1,2 (2)	2,3 (2)	–	–
5	3–7 (4)	2–5 (3)	2–5 (2)	5–9 (7)	4–7 (4)	3–7 (4)	4–7 (5)	4–7 (4)	–	–	–
6	1–3 (2)	1,2 (1)	1,2 (1)	1–3 (2)	1,2 (1)	1	1	1,2 (1)	–	–	–
7	2–4 (3)	2–5 (3)	2–5 (3)	2–8 (4)	2–5 (2)	2–6 (4)	1,3 (1)	1,2 (2)	–	–	–
8	1,2 (1)	–	–	2–4 (3)	2–4 (2)	1–3 (2)	1,3 (2)	1,2 (2)	–	–	–
9	2–4 (2)	1	1	1	1	1	1	1	8–13 (11)	–	–
10	1–3 (2)	–	1–3 (2)	2,3 (3)	1,2 (2)	1–3 (2)	1,2 (2)	2,3 (2)	–	–	–
11	1–5 (3)	–	0–2 (0)	1,2 (1)	1,2 (1)	1	1,2 (1)	1–4 (2)	–	–	–
12	2,3 (2)	–	–	–	–	–	–	–	–	–	–
13	0–2 (0)	–	–	–	–	–	–	–	–	–	–
14	–	–	–	–	1	1	1	1	1	–	–

Table 14. Number of branches for setae of fourth-instar larva of *An. mirans*: range (mode).

Seta no.	Head C	Thorax			Abdominal segments								
		P	M	T	I	II	III	IV	V	VI	VII	VIII	X
0	1	1	–	–	–	1	1	1	1	1	1	1	–
1	1	15–20 (15)	25–34 (28)	1–4 (3)	3–7 (3)	10–17 (10)	14–21 (17)	16–21 (18)	14–22 (18)	11–21 (17)	10–16 (13)	1	1–2 (1)
2	1	10–14 (12)	1–5 (3)	1	1–4 (3)	5–10 (7)	5–8 (7)	3–5 (4)	3–5 (4)	4–6 (5)	6–10 (10)	4–12 (7)	13–23
3	1–2 (1)	1	1	5–11 (7)	1–3 (2)	1	1–2 (1)	1–3 (2)	1	1	3–5 (3)	5–7 (7)	6–14 (10)
4	1–3 (1)	13–18 (13)	1–3 (2)	1–5 (3)	3–7 (5)	4–9 (7)	3–5 (4)	3–5 (4)	2–3 (3)	1–2 (1)	1	1	9
5	7–14 (10)	21–51 (33)	1	27–37 (27)	3–4 (3)	4–6 (6)	3–6 (3)	3–5 (3)	3–5 (3)	3–6 (4)	3–5 (4)	3–4 (3)	–
6	10–17 (13)	1	4–5 (4)	1–3 (2)	8–20 (8)	13–28 (19)	11–15 (15)	2–3 (3)	2–3 (3)	3–7 (3)	4–9 (5)	1–S	4–9 (5)
7	12–18 (15)	22–30	2–3 (3)	16–31 (1)	9–17 (1)	11–24 (17)	3–7 (6)	4–7 (5)	3–8 (4)	2–3 (3)	2–5 (3)	2–S	5–8 (8)
8	1–2 (2)	19–28 (28)	19–35 (24)	10–26 (0)	–	2–4 (2)	1–3 (2)	2–3 (2)	2–4 (3)	2–4 (3)	3–6 (5)	6–S	2–5 (3)
9	1–6 (2)	1	1	1–2 (1)	3–7 (5)	5–10 (8)	5–8 (7)	5–7 (5)	4–7 (5)	5–9 (6)	5–8 (6)	7–S	1–2 (1)
10	1–2 (1)	1–2 (1)	1–2 (1)	1	1	2–3 (3)	1	1–2 (1)	1–2 (1)	1–4 (4)	1–9 (7)	8–S	2–6 (4)
11	17–29 (27)	1	1	1	2–4 (3)	1	2–4 (3)	2–4 (3)	1–3 (3)	2–4 (3)	2–3 (2)	9–S	3–6 (3)
12	1–2 (2)	1–3 (1)	1	2–4 (2)	2–4 (3)	1	2–3 (3)	2–3 (3)	2–4 (3)	1–4 (1)	1–2 (1)	–	–
13	3–6 (3)	3–6 (5)	4–9 (7)	2–3 (3)	3–8 (5)	3–7 (5)	3–6 (4)	3–4 (3)	3–4 (3)	6–12 (9)	2–3 (2)	–	–
14	3–5 (5)	7–12 (8)	6–12 (9)	–	–	–	–	1	1	1	1	1	–
15	5–8 (6)	–	–	–	–	–	–	–	–	–	–	–	–

0.05–0.11 mm (mean 0.08 mm), 5-C longer than antenna, extending beyond anterior margin of head, with 7–14 branches, 6-C with 10–17 branches, 7-C with 12–18 branches. Thorax: tubercles of all large setae light to medium brown; seta 1-P with 15–20 branches, tubercle of seta 1-P usually joined basally to tubercle of 2,3-P, rarely separate, both tubercles with strong, pointed or somewhat rounded apical tooth projecting forward over base of 1,2-P, 14-P with 7–12 branches; 4-M single to triple, 6-M usually 4-branched (4,5), 14-M with 6–12 branches; 3-T moderately developed, not palmate, with 5–11 narrow branches arising from distinct level on stem. Abdomen: seta 2-I single to 4-branched, 3-I single to triple, 9-I with 3–7 branches; 1-II poorly developed, more pigmented than 1-I, with 10–17 leaflets arising from distinct level of a moderately pigmented stem, 13-IV with 3,4 branches, ratio of length of 13-IV to 10-IV 0.80–1.14 (mean 0.93); 1-VII with 10–16 moderately broad lanceolate leaflets with minute apicolateral serrations and slender apical filament; 1-X single or double; pecten teeth 13–15, with 5,6 long teeth alternating with 7–10 short teeth.

Material examined. HOLOTYPE: adult female with associated larval and pupal exuviae on microscope slide, acc. no. 316–1, collected by Yiau-Min Huang and Peyton, 18 July 1975, deposited in the NMNH. Type locality: SRI LANKA, Western, Kalatuara, Morapitiya, Sinharaja Forest Reserve (6°33'N 80°19'E). PARATYPES: INDIA, Karnataka (Mysore), Kârwâr, north Kanara (14°53'N 74°35'E), 3 September 1902, acc. no. 330, 1F; 1 October 1902, acc. no. 332, 1M; acc. no. 333, 1F; acc. no. 335, 1M; acc. no. 336, 1M; acc. no. 357, 10 October 1902, 1F; acc. no. 361, 12 October 1902, 1F; acc. no. 373, 29 October 1902, 1F; acc. no. 381, 12 January 1903, 1M. Madras (Tamil Nadu), Nilgiris, Buliar 11°25'N 76°30'E, 13 December 1977, acc. no. 655, 19F, 12M, 13Le, 29Pe; acc. no. 664, 23 December 1977, 4F, 4M, 4Le, 6Pe; acc. no. 676, 12 January 1978, 1F, 3M, 4Le, 4Pe. SRI LANKA, Central, Kandy, Udawatekelle Forest Reserve (7°15'N 80°37'E), June 1975, acc. no. 43, 1M, 1Le, 1Pe, all collected by Peyton and Huang; Kandy, Wakarawatta, Roseneath (7°17'N 80°38'E), 25 March 1971, 1L, collected by BA Harrison *et al.* Sabaragamuwa, Ratnapura, Vaddagala, Sinharaja Forest Reserve (6°24'N 80°27'E), acc. no. 286, 17 July 1975, 1M, 1Le, 1Pe; acc. no. 295, 17 July 1995, 9F, 15M, 19Le, 23Pe. Southern, Galle, Kanneliya Forest Reserve (6°14'N 80°20'E), 9 July 1975, acc. no. 199, 1F, 1Le, 1Pe. Western, Colombo, Labugama Reservoir (6°50'N 80°10'E), acc. no. 422, 7 August 1975, 2F, 4M, 4Le, 4Pe; Kalatuara, Morapitiya, Sinharaja Forest Reserve (6°33'N 80°19'E), 18 July 1975, acc. no. 316, 4F, 5M, 4Le, 4Pe, all collected by Huang and Peyton. Paratypes are deposited in the NMNH. Total material of *An. mirans* examined: 1049 specimens comprising 235F, 183M, 210Le, 359Pe, 63L derived from 94 separate collections from natural habitats (14 adult, 80 immature).

Distribution. *Anopheles mirans* is known from south-western India (States of Kerala, Karnataka, Tamil Nadu)

and Sri Lanka (Central, Sabaragamuwa, Southern, North-western and Western Provinces).

Bionomics. In Tamil Nadu, India, immatures of *An. mirans* were found in shallow, large and small ground pools, shallow or deep large or small rock pools, flood pools and swamps. The water was clear or turbid, in partial or heavy shade, sometimes with decaying leaves. The larval habitats were situated in mountainous areas.

In Sri Lanka, *An. mirans* immatures were collected in cultivated areas, in ditches along road sides, animal footprints, concrete wells, small rock pools, wheel ruts, ground pools, flood pools, pools along side roads, small or large rock pools, swamps, rock pools along stream edges, and in a semipermanent gem pit. The water was always fresh, stagnant, coloured or turbid or clear, in partial or heavy shade and usually with leaves and sticks. The habitats were in secondary rain forest with mixed scrub vegetation, situated in mountainous terrain, at elevations from 200 to 900 m above sea level. Immatures of *An. mirans* were found in association with *Cx. bailyi* Barraud, *Cx. fragilis* Ludlow, *Cx. lasiopalis* Sirivanakarn, *Cx. mimulus*, *Cx. minutissimus* (Theobald), *Cx. pallidothorax* Theobald, *Cx. quadripalpis* (Edwards), *Cx. uniformis* (Theobald), *Cx. wardi* Sirivanakarn, *Ur. bicolor* and *An. maculatus*.

Adults of *An. mirans* were collected at Kandy, Central Province of Sri Lanka, in a small forest (104 ha) bordering the city in central hills of the country at elevation of 518 m above sea level. The vegetation consisted of tall trees with dense undergrowth. Larval habitats were muddy water in car tracks, tyre marks and similar depressions on an infrequently used gravel road. The water was clean except for a few fallen leaves, and it was heavily shaded. Additionally, immatures were found in association with several other Culicinae species but were the only anopheline larva encountered throughout the rainy season. However, when the drought was approaching, immatures of *An. mirans* declined in numbers (Mendis *et al.*, 1984).

Wild-caught females of *An. mirans*, from a small locality in foothills of the eastern Nilgiris, were found infected with sporozoites and oocysts of *P. cynomolgi* Mayer and *P. inui* Halberstaedter & von Prowazek (Choudhury *et al.*, 1963b). Additionally, this species was implicated as a natural vector of simian malaria parasites in Sri Lanka where adults were found naturally infected with sporozoites of *P. inui shortii* Bray and *P. fragile* Dissanaiké, Nelson & Garnham (Nelson *et al.*, 1971). In another study, under laboratory conditions, females of *An. mirans* became infected with simian malarias (*P. cynomolgi* and *P. inui*) when fed on *Macaca radiata* (Saint-Hilaire), but further transmission by these *An. mirans* was not attempted (Choudhury *et al.*, 1963a).

Systematics. *Anopheles mirans* has been mostly misidentified as *An. elegans* since Cogill (1903). *Anopheles mirans* belong to the Hackeri Subgroup based on the proboscis being noticeably longer than the forefemur. Possible due to its restricted distribution and relative non-availability, this species has received less attention

than other members of the *Leucosphyrus* Group. Adult females can be recognized by the following combination of characters: (1) proboscis longer than forefemur (ratio between proboscis length and forefemur length 1.19–1.31); (2) proboscis longer than maxillary palpus, ratio between palpus length and proboscis length 0.84–0.93; (3) proboscis entirely dark-scaled; (4) PSD spot of vein R without pale interruptions; (5) number of pale interruptions on PSD-PD spots of vein R always 2; (5) ratio of length of apical pale band of palpomere 5 to length of basal dark band of palpomere 5 0.83–2.67; and (6) apical pale band of hindtibia with continuous dark longitudinal stripe extending into basal portion.

Fourth-instar larvae of *An. mirans* can be recognized by the following combination of characters: (1) seta 5-C conspicuously longer than antenna; (2) basal tubercle of seta 1-P with prominent tooth or spine on posterodorsal margin; (3) seta 4-C long, extending noticeably beyond base of 2-C; (4) seta 1-II poorly developed with very narrow leaflets; (5) ratio of length of seta 13-IV to 10-IV 0.80–1.14 (mean = 0.93); (6) seta 1-II with 10–17 branches; (7) seta 13-IV,V with 3,4 branches; (8) seta 2-VII with 6–10 branches; and (9) seta 9-VII with 5–8 branches. The pupal stage of *An. mirans* cannot be distinguished from other members of the Hackeri Subgroup.

Discussion

With their beautiful adult scale-patterns of fine white spots and dark background facies, all species of the *Leucosphyrus* Group are readily recognizable from their distinctive broad white tibio-tarsal band on the hind-legs. As they transmit the bulk of forest malaria in oriental countries, these mosquitoes are much feared for the extremely high vectorial capacity of some but not all species, notably the aptly named *An. dirus* – meaning danger. Due to the variability of their elaborate ornamentation, however, separating the many species of this group will always be morphologically challenging. As with other groups of anopheline mosquitoes, therefore, taxonomic progress at the species level has depended increasingly on genetic criteria and specific micro-morphological characters on the immature stages. Combinations of such evidence have allowed us to define the six new species morphologically described and formally named here. They fit the expected criteria (see Fig. 1 in Peyton, 1990) for belonging to the Hackeri Subgroup or the *Leucosphyrus* Subgroup, based on length ratios of the proboscis/forefemur and the maxillary palpus/proboscis. Thus, we are preparing a taxonomic revision of the complete group (Sallum *et al.*, 2005) including 20 species. This should improve the prospects for understanding their bioecology and specific vector roles, and facilitate better targeting of control operations. Studies on distribution, seasonality, biting activity, feeding habits, longevity, infectivity, vector competence and vectorial capacity depend on the correct identification of such species in each locality. Fortunately it is now possible to distinguish between these species by

genetic and molecular procedures, augmented by the morphological characters we have determined.

In the light of these six new species, we have to re-examine the differential distributions and specific bionomics of typical *An. balabacensis*, *An. dirus*, *An. leucosphyrus* and other members of the *Leucosphyrus* Group, particularly to understand their roles as vectors of malaria (*Plasmodium* spp.) and filaria (*Wuchereria* spp.) parasites among primates in and near forests. For example, *An. latens* (which has been largely misidentified as *An. leucosphyrus* s.s.) is anthropophilic and apparently serves as an important vector of human malaria in villages and forested areas of Sarawak, East Malaysia (Colless, 1956; Zulueta, 1956; Chang *et al.*, 1995), and of South Kalimantan, Indonesia (White, 1983; Harbach *et al.*, 1987). *Anopheles latens* is also responsible for most transmission of *Wuchereria bancrofti* to humans in Sarawak (Zulueta & LaChance, 1956; Zulueta, 1957), whereas *Wuchereria kalimantani* of leaf monkeys appears to be transmitted by *An. balabacensis* in Kalimantan (Atmosoedjono *et al.*, 1993). Even so, *An. balabacensis* is a major vector of malaria in eastern Borneo and adjacent islands.

Anopheles baimaii is evidently another important vector of *P. vivax* and *P. falciparum* in forest, forest fringes and foothill areas throughout its wide range (from India to China), and its vectorial capacity seems to be comparable to that of *An. dirus* in Thailand. Records show that it is vector in Bangladesh (Rahman *et al.*, 1977; Rosenberg & Maheswary, 1982), Burma (Khin-Maung-Kyi & Winn, 1976; Tun-Lin *et al.*, 1995) and north-east India (Dutta *et al.*, 1991; Prakash *et al.*, 2001, 2005). Assessment of the vector role of *An. baimaii* in Thailand is more complex since it occurs in sympatry with other species of the complex (Baimai *et al.*, 1988c), combining their contributions to the dynamics of malaria transmission. However, *An. baimaii* seems to have a more western distribution, being the predominant member of the *Leucosphyrus* Group and an efficient vector of human *Plasmodium* in western Thailand (Baimai *et al.*, 1988c). Despite being mainly a forest mosquito, *An. baimaii* withstands anthropic environmental changes. In Myanmar, peridomestic populations of *An. baimaii* breed in wells (Tun-Lin *et al.*, 1987; Oo *et al.*, 2003) and have different esterase isozyme profiles from those of forested populations (Tun-Lin *et al.*, 1988).

Anopheles cracens is another strongly anthropophilic species (Baimai *et al.*, 1988c) that may also contribute to the transmission of both malaria and filaria parasites. Additionally, *An. cracens* is competent to transmit *Plasmodium cynomolgi* (Klein *et al.*, 1991) and *Brugia pahangi* (Zahedi & White, 1994). The other man-biting species such as *An. elegans* in India and *An. scanloni* in Thailand should be checked in relation to the epidemiology of both malaria and filariasis, since they have not been identifiable hitherto, so their vectorial importance may have been underestimated. The non-anthropophilic *Anopheles mirans* may be a natural vector of *P. inui shortii* and *P. fragile* in Sri Lanka (Nelson *et al.*, 1971) as well as *P. cynomolgi* and *P. inui* in foothills of south India (Choudhury *et al.*,

1963b). As a laboratory vector, *An. mirans* was readily infected with *P. cynomolgi* and *P. inui* from *Macaca radiata* (Choudhury *et al.*, 1963a).

Under natural forest conditions, most if not all members of the Leucosphyrus Group apparently feed primarily on monkeys in the tree-tops, transmitting various plasmodia among them. Two species of simian malaria have been detected in *An. dirus* s.l. (Cheong *et al.*, 1965), five species in *An. hackeri* and three species in *An. introlatus*, and *An. balabacensis*, *An. puyutensis* and *An. leucosphyrus* s.l. have also been implicated (Warren & Wharton, 1963; Coatney *et al.*, 1971). Presumably all the oriental primate malarias are transmitted to some extent by mosquitoes of the Leucosphyrus Group. For example, the *Macaca* parasites *Plasmodium knowlesi* and possibly *P. coatneyi* and *P. inui* have been detected in *An. balabacensis* from Palawan Island of The Philippines (Tsukamoto *et al.*, 1978), although *P. cynomolgi* can be harmful to *An. dirus* s.l. under laboratory conditions (Klein *et al.*, 1982). Circumstantial evidence in Sabah led to proof of *An. balabacensis* vector competence for *P. pitheci* and *P. silvaticum* from orang-utans (Peters *et al.*, 1976). Hence, the need to clarify the veterinary as well as medical importance of all species in the Leucosphyrus Group.

Acknowledgements

This research was performed under a Memorandum of Understanding between the Walter Reed Army Institute of Research and the Smithsonian Institution, with institutional support provided by both organizations. This information reflects the views of the authors and should be not construed to represent those of the Department of the Army or the Department of Defense. This project was completed while M.A.M.S. was senior research fellow supported by the U.S. National Research Council (NRC) Grant DAMD 17-00-2-0002. Sadly, E. L. Peyton passed away before this paper was completed, having done so much to facilitate our accomplishment of this study.

We are most grateful to Rampa Rattanarithikul (AFRIMS), Visut Baimai, Shiva Ramalingam and the late Chris Green, Mahidol University, Bangkok, and B. Harrison (DENR Public Health Pest Management North Carolina) for their insight and cooperation over species diagnosis. Wayne Mathis (Department of Entomology, NMNH, Smithsonian Institution, Washington D.C.) and Christian Thompson (Systematic Entomology Laboratory, U. S. Department of Agriculture, and Department of Entomology, NMNH, Smithsonian Institution) are acknowledged for providing critical comments that fostered the solution of the nomenclature regarding *An. elegans*. Ralph E. Harbach (Department of Entomology, BMNH, London) is acknowledged for his critical review and valuable suggestions that greatly improved the manuscript. Most importantly we thank all colleagues, many anonymous as well as those mentioned in the text, who collected specimens used in this study, particularly those at the

AFRIMS, Bangkok, and at the Institute of Medical Research, Kuala Lumpur.

References

- Atmosoedjono, S., Djoharti, Purnomo & Bangs, M.J. (1993) *Anopheles balabacensis* (Diptera: Culicidae), a vector of *Wuchereria kalimantani* (Nematoda: Onchocercidae) in East Kalimantan (Borneo), Indonesia. *Medical and Veterinary Entomology*, **7**, 390–392.
- Audtho, M., Tassanakajon, A., Boonsaeng, V., Tpiankijaguin, S. & Panyim, S. (1995) Simple nonradioactive DNA hybridization method for identification of sibling species of *Anopheles dirus* (Diptera: Culicidae) complex. *Journal of Medical Entomology*, **32**, 107–111.
- Baimai, V. (1988a) Constitutive heterochromatin differentiation and evolutionary divergence of karyotype in Oriental *Anopheles* (Cellia). *Pacific Science*, **42**, 13–27.
- Baimai, V. (1988b) Population cytogenetics of the malaria vector *Anopheles leucosphyrus* group. *Southeast Asian Journal of Tropical Medicine and Public Health*, **19**, 667–680.
- Baimai, V., Andre, R.G., Harrison, B.A., Kijchalao, U. & Panthusiri, L. (1987) Crossing and chromosomal evidence for two additional sibling species within the taxon *Anopheles dirus* Peyton and Harrison (Diptera: Culicidae) in Thailand. *Proceedings of the Entomological Society of Washington*, **89**, 157–166.
- Baimai, V., Green, C.A., Andre, R.G., Harrison, B.A. & Peyton, E.L. (1984) Cytogenetic studies of some species complexes of *Anopheles* in Thailand and Southeast Asia. *Southeast Asian Journal of Tropical Medicine and Public Health*, **15**, 536–546.
- Baimai, V., Harbach, R.E. & Kijchalao, U. (1988a) Cytogenetic evidence for a fifth species within the taxon *Anopheles dirus* in Thailand. *Journal of the American Mosquito Control Association*, **4**, 333–338.
- Baimai, V., Harbach, R.E. & Sukowati, S. (1988b) Cytogenetic evidence for two species within the current concept of the malaria vector *Anopheles leucosphyrus* in Southeast Asia. *Journal of the American Mosquito Control Association*, **4**, 44–50.
- Baimai, V., Harrison, B.A. & Somchit, L. (1981) Karyotype differentiation of three anopheline taxa in the Balabacensis Complex of Southeast Asia (Diptera: Culicidae). *Genetica*, **57**, 81–86.
- Baimai, V., Kijchalao, U., Sawadwongporn, P. & Green, C.A. (1988c) Geographic distribution and biting behaviour of four species of the *Anopheles dirus* complex (Diptera: Culicidae) in Thailand. *Southeast Asian Journal of Tropical Medicine and Public Health*, **19**, 151–161.
- Baimai, V., Poopittayasataporn, A. & Kijchalao, U. (1988d) Cytological differences and chromosomal rearrangements in four members of the *Anopheles dirus* complex (Diptera: Culicidae). *Genome*, **30**, 372–379.
- Baimai, V., Thu, M.M., Paing, M. & Maheswary, N.P. (1988e) Distribution and chromosomal polymorphism of the malaria vector *Anopheles dirus* species D. *Southeast Asian Journal of Tropical Medicine and Public Health*, **19**, 661–665.
- Baimai, V. & Traipakvasin, A. (1987) Intraspecific variation in sex heterochromatin of species B of the *Anopheles dirus* complex in Thailand. *Genome*, **29**, 401–404.
- Belkin, J.N. (1962) *The Mosquitoes of the South Pacific* (Diptera, Culicidae), Vol. 1. University of California Press, Los Angeles.
- Bhat, H.R. (1988) A note on *Anopheles dirus* Peyton and Harrison, 1979 [*An. balabacensis* (sensu lato) Baisas, 1936] in India. *Indian Journal of Malariology*, **25**, 103–105.

- Chang, M.S., Doraisingam, P., Hardin, S. & Nagum, N. (1995) Malaria and filariasis transmission in a village/forest setting in Baram District, Sarawak, Malaysia. *Journal of Tropical Medicine and Hygiene*, **98**, 192–198.
- Chareonviriyaphap, T., Bangs, M.J. & Ratanatham, S. (2000) Status of malaria in Thailand. *Southeast Asian Journal of Tropical Medicine and Public Health*, **31**, 225–237.
- Cheong, W.H., Warren, McW., Omar, A.H. & Mahadevan, S. (1965) *Anopheles balabacensis* identified as a vector of simian malaria in Malaysia. *Science*, **150**, 1314–1315.
- Choudhury, D.S., Mohan, B.N., Prakash, S. & Ramakrishnan, S.P. (1963a) Experimental susceptibility of anopheline mosquitoes to simian malaria in the Nilgiris, Madras State, South India. *Indian Journal of Malariology*, **17**, 237–242.
- Choudhury, D.S., Wattal, B.L. & Ramakrishnan, S.P. (1963b) Incrimination of *Anopheles elegans* James (1903) as a natural vector of simian malaria in the Nilgiris, Madras State, India. *Indian Journal of Malariology*, **17**, 243–247.
- Christophers, S.R. (1924) Provisional list and reference catalogue of the Anophelini. *Indian Medical Research Memoir*, **3**, 1–105.
- Christophers, S.R. (1933) *The Fauna of British India, Including Ceylon and Burma Diptera*, Vol. IV, Family Culicidae Tribe Anophelini. Taylor & Francis, London.
- Coatney, G.R., Collins, W.E., Warren, McW. & Contacos, P.G. (1971) *The Primate Malariae*. National Institutes of Health. U.S. Department of Health, Education and Welfare, Bethesda, Maryland.
- Cogill, H. (1903) The *Anopheles* of Karwar (North Kanara). *Journal of the Bombay Natural History Society*, **15**, 327–336.
- Colless, D.H. (1956) The *Anopheles leucosphyrus* group. *The Transactions of the Royal Entomological Society of London*, **108**, 37–116.
- Colless, D.H. (1957) Further notes on the systematics of the *Anopheles leucosphyrus* group (Diptera: Culicidae). *Proceedings of the Entomological Society of London (B)*, **26**, 131–139.
- Damrongphol, P. & Baimai, V. (1989) Scanning electron microscope observations and differentiation of eggs of the *Anopheles dirus* complex. *Journal of the American Mosquito Control Association*, **5**, 563–568.
- Daniels, G.W. (1908) Breeding grounds of Culicidae. *Studies from Institute for Medical Research, Federated Malay States*, **3**, 1–7.
- Dutta, P., Bhattacharyya, D.R. & Dutta, L.P. (1989a) Incrimination of *Anopheles dirus* as a vector of malaria in Dibrugarh district, Assam. *Indian Journal of Malariology*, **26**, 149–152.
- Dutta, P., Bhattacharyya, D.R. & Dutta, L.P. (1991) Epidemiological observations on malaria in some parts of Tengakhat PHC Dibrugarh District Assam. *Indian Journal of Malariology*, **28**, 121–128.
- Dutta, P., Bhattacharyya, D.R., Khan, S.A., Sharma, C.K. & Mahanta, J. (1996) Feeding patterns of *Anopheles dirus*, the major vector of forest malaria in northeast India. *Southeast Asian Journal of Tropical Medicine and Public Health*, **27**, 378–381.
- Dutta, P., Bhattacharyya, D.R., Sharma, C.K. & Dutta, L.P. (1989b) The importance of *Anopheles dirus* (*A. balabacensis*) as a vector of malaria in northeast India. *Indian Journal of Malariology*, **26**, 95–101.
- Dutta, P., Bhattacharyya, D.R., Sharma, C.K. & Dutta, L.P. (1992) Anopheline fauna of parts of Tirap district, Arunachal Pradesh with reference to malaria transmission. *Indian Journal of Medical Research Section A*, **95**, 245–249.
- Dutta, P., Dev, V. & Bhattacharyya, D.R. (1993) Anopheline fauna and malaria incidence in Changlang District (Arunachal Pradesh). *Indian Journal of Malariology*, **30**, 135–143.
- Dutta, P., Khan, S.A., Sharma, C.K. & Mahanta, J. (1997) A report of mosquito fauna survey and vector incrimination in Goalpara district of Assam. *Indian Journal of Malariology*, **34**, 204–207.
- Green, C.A., Munstermann, L.E., Tan, S.G., Panyim, S. & Baimai, V. (1992) Population genetic evidences for species A, B, C, and D of the *Anopheles dirus* complex in Thailand and enzyme electromorphs for their identification. *Medical and Veterinary Entomology*, **6**, 29–36.
- Green, C.A., Rattanarithikul, R., Pongparit, S., Sawadwongporn, P. & Baimai, V. (1991) A newly-recognized vector of human malarial parasites in the Oriental region *Anopheles* (*Cellia*) *pseudowillmori* (Theobald, 1910). *Transactions of the Royal Society of Tropical Medicine and Hygiene*, **85**, 35–36.
- Harbach, R.E. (2004) The classification of genus *Anopheles* (Diptera: Culicidae): a working hypothesis of phylogenetic relationships. *Bulletin of Entomological Research*, **94**, 537–553.
- Harbach, R.E., Baimai, V. & Sukowati, S. (1987) Some observations on sympatric populations of the malaria vectors *Anopheles leucosphyrus* and *Anopheles balabacensis* in a village-forest setting in South Kalimantan. *Southeast Asian Journal of Tropical Medicine and Public Health*, **18**, 241–247.
- Harbach, R.E. & Knight, K.L. (1980) *Taxonomists' Glossary of Mosquito Anatomy*. Plexus Publishing, Inc. Marlton, New Jersey.
- Harbach, R.E. & Knight, K.L. (1981) Corrections and additions to Taxonomists' Glossary of Mosquito Anatomy. *Mosquito Systematics*, **13**, 201–217.
- Hii, J.L.K. (1982) *Laboratory studies on three member species of the Anopheles balabacensis complex (Diptera: Culicidae)*. PhD Thesis, University of London.
- Hii, J.L.K. (1984a) Insecticide susceptibility of three cryptic species of the *Anopheles balabacensis* complex. *Southeast Asian Journal of Tropical Medicine and Public Health*, **15**, 104–111.
- Hii, J.L.K. (1984b) Involvement of the X-chromosome in hybrid male sterility from crosses between species A and species B of the taxon *Anopheles dirus*. *Mosquito News*, **44**, 192–196.
- Hii, J.L.K. (1985) Genetic investigations of laboratory stocks of the complex of *Anopheles balabacensis* Baisas (Diptera: Culicidae). *Bulletin of Entomological Research*, **75**, 185–197.
- Hii, J.L.K. (1986) Multivariate morphometrics of the immature stages of the *Anopheles balabacensis* complex (Diptera: Culicidae). *Mosquito Systematic*, **18**, 125–133.
- Huong, N.T., Sonthayanon, P., Ketterman, A.J. & Panyim, S. (2001) A rapid polymerase chain reaction based method for identification of the *Anopheles dirus* sibling species. *Southeast Asian Journal of Tropical Medicine and Public Health*, **32**, 615–620.
- ICZN (1999) *International Code of Zoological Nomenclature*, 4th edn. The International Trust for Zoological Nomenclature, The Natural History Museum, London.
- James, S.P. & Liston, W.G. (1904) *A Monograph of the Anopheles Mosquitoes of India*. Thacker, Spink, Calcutta.
- James, S.P. & Liston, W.G. (1911) *Anopheline Mosquitoes of India*, 2nd edn. Thacker, Spink, Calcutta.
- James, S.P. & Stanton, A.T. (1912) Revision of the names of Malayan anophelines. *Paludism*, **5**, 59–63.
- Kanda, T., Takai, K., Chiang, G.L., Cheong, W.H. & Sucharit, S. (1981) Hybridization and some biological factors of seven strains of the *Anopheles leucosphyrus* group (Reid, 1968). *Japanese Journal of Sanitary Zoology*, **32**, 321–329.
- Kanda, T., Takai, K., Chiang, G.L., Loong, K.P., Sucharit, S. & Cheong, W.H. (1983) Phylogenetic interpretation and chromosomal polymorphism among nine strains of human malaria vectors of the *Anopheles leucosphyrus* group. *Japanese Journal of Genetics*, **58**, 193–208.

- Khan, S.A., Handique, R., Tewari, S.C., Dutta, P., Narain, K. & Mahanta, J. (1998) Larval ecology and mosquito fauna of Upper Brahmaputra valley, northeast India. *Indian Journal of Malariology*, **35**, 131–145.
- Khin-Maung-Kyi & Winn, S.M. (1976) *Studies on Malaria and Anopheles balabacensis balabacensis and Anopheles minimus in Burma*. Unpublished document. WHO/Mal/76.875. World Health Organization, Geneva.
- Klein, T.A., Harrison, B.A., Andre, R.G., Whitmire, R.E. & Inlao, I. (1982) Detrimental effect of *Plasmodium cynomolgi* infections on the longevity of *Anopheles dirus*. *Mosquito News*, **42**, 265–271.
- Klein, T.A., Harrison, B.A., Dixon, S.V. & Burge, J.R. (1991) Comparative susceptibility of Southeast Asian *Anopheles* mosquitoes to the simian malaria parasite *Plasmodium cynomolgi*. *Journal of the American Mosquito Control Association*, **7**, 481–487.
- Leicester, G.F. (1903) A breeding place of certain forest mosquitoes in Malaya. *Journal of Tropical Medicine*, **6**, 291–292.
- Leicester, G.F. (1908a) Report of an investigation of the mosquitoes of Port Swettenham, Klang and Jugra. *Studies of the Institute of Medical Research of the Federated Malay States*, **3**, 7–14.
- Leicester, G.F. (1908b) The Culicidae of Malaya. *Studies of the Institute of Medical Research of the Federated Malay States*, **3**, 18–261.
- Leicester, G.F. (1908c) Notes as to the pathogenicity of certain mosquitoes with respect to filaria nocturna and malignant tertian malaria. *Studies of the Institute of Medical Research of the Federated Malay States*, **3**, 267–269.
- Macan, T.T. (1950) Malaria survey of the Arakan region of Bengal and Burma. *Parasitology*, **40**, 290–297.
- Manguin, S., Kengne, P., Sonnier, L., Harbach, R.E., Baimai, V., Trung, H.D. & Coosemans, M. (2002) SCAR markers and multiplex PCR-based identification of isomorphic species in the *Anopheles dirus* complex in Southeast Asia. *Medical and Veterinary Entomology*, **16**, 46–54.
- May, K. (1987) Karyological identification of two taxa of the *Anopheles balabacensis* complex from Burma. *Genome*, **29**, 886–887.
- Mendis, K., Ithalamulla, R.L., Peyton, E.L. & Nanayakkara, S. (1984) Biology and descriptions of the larva and pupa of *Anopheles (Cellia) elegans* James (1903). *Mosquito Systematics*, **1983** (15), 318–324.
- Misra, S.P., Nandi, J., Narasimham, M.V.V.L. & Rajagopal, R. (1993) Malaria transmission in Nagaland, India: Part I. Anophelines and their seasonality. *Journal of Communicable Diseases*, **25**, 62–66.
- Moore, S.J., Zunwei, D., Hongning, Z., Xuezhong, W., Hongbing, L., Yujiang, X. & Hill, N. (2001) The efficacy of different mosquito trapping methods in a forest-fringe village, Yunnan Province, southern China. *Southeast Asian Journal of Tropical Medicine and Public Health*, **32**, 282–289.
- Nandi, J., Kaul, S.M., Sharma, S.N. & Lal, S. (2000) Anthropophily of anophelines in Duars of West Bengal and other regions of India. *Journal of Communicable Diseases*, **32**, 95–99.
- Nandi, J., Misra, S.P., Rajagopal, R. & Narasimham, M.V.V.L. (1993) Present perspectives of malaria transmission in Boko Area of Assam. *Journal of Communicable Diseases*, **25**, 18–26.
- Nandi, J., Rao, J.S., Dasgupta, R.K. & Sharma, R.S. (1996) Ecological observations on the anopheline mosquitoes of Jalpaiguri duars, West Bengal. *Journal of Communicable Diseases*, **28**, 279–286.
- Nelson, P., Jayasuriya, J.M.R. & Bandarawatte, B.V.P.C. (1971) The establishment of *Anopheles elegans* as a natural vector of simian malaria in Ceylon. *Ceylon Journal of Medical Science*, **20**, 46–51.
- Oo, T.T., Storch, V. & Becker, N. (2003) *Anopheles dirus* and its role in malaria transmission in Myanmar. *Journal of Vector Ecology*, **28**, 175–183.
- Panyim, S., Yasothornsrikul, S., Tungpradubkul, S., Baimai, V., Rosenberg, R., Andre, R.G. & Green, C.A. (1988) Identification of isomorphic malaria vectors using a DNA probe. *American Journal of Tropical Medicine and Hygiene*, **38**, 47–49.
- Peters, W., Garnham, P.C.C., Killick-Kendrick, R., Rajapaksa, N., Cheong, W.H. & Cadigan, F.C. (1976) Malaria of the orangutan (*Pongo pygmaeus*) in Borneo. *Philosophical Transactions of the Royal Society of London, B. Biological Sciences*, **275**, 439–482.
- Peyton, E.L. (1990) A new classification for the Leucosphyrus Group of *Anopheles (Cellia)*. *Mosquito Systematics*, **21** [1989], 197–205.
- Peyton, E.L. & Harrison, B.A. (1979) *Anopheles (Cellia) dirus*, a new species of the Leucosphyrus Group from Thailand (Diptera: Culicidae). *Mosquito Systematics*, **11**, 40–52.
- Peyton, E.L. & Harrison, B.A. (1980) *Anopheles (Cellia) takasagoensis* Morishita 1946, an additional species in the Balabacensis Complex of Southeast Asia (Diptera: Culicidae). *Mosquito Systematics*, **12**, 335–347.
- Peyton, E.L. & Ramalingam, S. (1988) *Anopheles (Cellia) nemophilous*, a new species of the Leucosphyrus Group from peninsular Malaysia and Thailand (Diptera: Culicidae). *Mosquito Systematics*, **20**, 272–299.
- Poopittayasataporn, A. & Baimai, V. (1995) Polytene chromosome relationships of five species of the *Anopheles dirus* complex in Thailand. *Genome*, **38**, 426–434.
- Prakash, A., Bhattacharya, D.R., Mohapatra, P.K. & Mahanta, J. (1998a) Insecticide susceptibility of *Anopheles dirus* in Assam. *Journal of Communicable Diseases*, **30**, 291–292.
- Prakash, A., Bhattacharya, D.R., Mohapatra, P.K. & Mahanta, J. (1997a) Indoor biting behaviour of *Anopheles dirus* Peyton and Harrison, 1979 in upper Assam, India. *Mosquito Borne Diseases Bulletin*, **14**, 31–37.
- Prakash, A., Bhattacharya, D.R., Mohapatra, P.K. & Mahanta, J. (1997b) Seasonal prevalence of *Anopheles dirus* and malaria transmission in a forest fringed village of Assam, India. *Indian Journal of Malariology*, **34**, 117–125.
- Prakash, A., Bhattacharya, D.R., Mohapatra, P.K. & Mahanta, J. (1997c) Breeding and day resting habitats of *Anopheles dirus* in Assam, India. *Southeast Asian Journal of Tropical Medicine and Public Health*, **28**, 610–614.
- Prakash, A., Bhattacharya, D.R., Mohapatra, P.K. & Mahanta, J. (1998b) House frequenting and host seeking mosquitoes in a forest fringed village of district Dibrugarh, Assam. *Entomon*, **23**, 191–195.
- Prakash, A., Bhattacharya, D.R., Mohapatra, P.K. & Mahanta, J. (2000a) Preliminary observations on man-mosquito contact in Soraipung village of District Dibrugarh, Assam (India). *Indian Journal of Malariology*, **37**, 97–102.
- Prakash, A., Bhattacharya, D.R., Mohapatra, P.K. & Mahanta, J. (2001) Estimation of vectorial capacity of *Anopheles dirus* in (Diptera: Culicidae) in a forested-fringed village of Assam (India). *Vector Borne Zoonotic Diseases*, **1**, 231–237.
- Prakash, A., Bhattacharya, D.R., Mohapatra, P.K. & Mahanta, J. (2002) Physico-chemical characteristics of breeding habitats of *Anopheles dirus* (Diptera: Culicidae) in Assam, India. *Journal of Environmental Biology*, **23**, 95–100.
- Prakash, A., Bhattacharya, D.R., Mohapatra, P.K. & Mahanta, J. (2005) Risk of malaria transmission by *Anopheles baimai* at different hours of night in Assam forest village. *Medical and Veterinary Entomology*, **19**, in press.
- Prakash, A., Mohapatra, P.K., Bhattacharya, D.R., Sharma, C.K., Goswami, B.K., Hazarika, N.C. & Mahanta, J. (2000b)

- Epidemiology of malaria outbreak (April/May, 1999) in Titabor Primary Health Centre, district Jorhat (Assam). *Indian Journal of Medical Research*, 111, 121–126.
- Prapanthadara, L., Promtet, N., Koottathap, S., Somboon, P. & Ketterman, A.J. (2000) Isoenzymes of glutathione S-transferase from the mosquito *Anopheles dirus* species B: the purification, partial characterization and interaction with various insecticides. *Insect Biochemistry and Molecular Biology*, 30, 395–403.
- Rahman, M., Elias, M.D. & Ameen, M. (1977) Bionomics of *Anopheles balabacensis balabacensis* Baisas (Diptera: Culicidae) in Bangladesh and its relation to malaria. *Bangladesh Journal of Zoology*, 5, 1–23.
- Ramsay, G.C. (1930) Some findings and observations in an anopheline malaria infectivity survey carried out in the Cachar District of Assam. *Indian Journal of Medical Research*, 18, 533–552.
- Rattanakul, R., Green, R.C., Panyim, S., Noigamol, C., Chanaimong, S. & Mahapibul, P. (1995) Larval habitats of malaria vectors around a transmission focus in northwestern Thailand. *Journal of the American Mosquito Control Association*, 11, 428–432.
- Reid, J.A. (1949) A preliminary account of the forms of *Anopheles leucosphyrus* Dönitz (Diptera: Culicidae). *Proceedings of the Royal Entomological Society of London B*, 18, 42–53.
- Reid, J.A. (1968) *Anopheles* mosquitoes of Malaya and Borneo. *Studies of the Institute of Medical Research of Malaya*, 31, 1–520.
- Reid, J.A. (1970) Systematics of malaria vectors. Anopheline systematics and malaria control, with special reference to Southeast Asia. *Miscellaneous Publication of the Entomological Society of America*, 7, 56–62.
- Rosenberg, R. (1982) Forest malaria in Bangladesh. III. Breeding habits of *Anopheles dirus*. *American Journal of the Tropical Medicine and Hygiene*, 31, 192–201.
- Rosenberg, R. & Maheswary, N.P. (1982) Forest malaria in Bangladesh. II. Transmission by *Anopheles dirus*. *American Journal of the Tropical Medicine and Hygiene*, 31, 183–191.
- Sawadipani, Y., Baimai, V. & Harrison, B.A. (1990) *Anopheles dirus* species E: chromosomal and crossing evidence for another member of the Dirus Complex. *Journal of the American Mosquito Control Association*, 6, 477–481.
- Seng, C.-M., Matusop, A. & Sen, F.-K. (1999) Differences in *Anopheles* composition and malaria transmission in the village settlements and cultivated farming zone in Sarawak, Malaysia. *Southeast Asian Journal of Tropical Medicine and Public Health*, 30, 454–459.
- Service, M.W. (1993) *Mosquito Ecology: Field Sampling Methods*. 2nd edn. Elsevier, Science Publishers, Essex, U.K.
- Sithiprasasna, R., Luepromchai, E. & Linthicum, K.J. (1996) Effects of sublethal dosages of methoprene on *Anopheles dirus* species A and B. *Journal of the American Mosquito Control Association*, 12, 483–486.
- Sucharit, S. & Choochote, W. (1983) Comparative studies on the morphometry of male genitalia and frequency of clasper movements during induced copulation of *Anopheles balabacensis* (Perlis Form) and *Anopheles dirus* (Bangkok colony strain). *Mosquito Systematics*, 15, 90–98.
- Swellengrebel, N.H. (1916) De anophelen van Nederlandsch Oost-Indië. Koloniaal Instituut te Amsterdam. Mededeeling, VII. *Afdeeling Tropische Hygiene*, 3, 1–182.
- Takai, K. (1986) Discrimination among strains of the mosquito *Anopheles leucosphyrus* group by electrophoresis of proteins. *Japanese Journal of Genetics*, 61, 45–59.
- Takai, K., Kanda, T., Cheong, W.H. & Loong, K.P. (1984) A recessive mutant, pale eye, and a cytoplasmic deleterious effect on viability in *Anopheles leucosphyrus* (Diptera: Culicidae). *Journal of Medical Entomology*, 21, 145–148.
- Tewari, S.C., Hiriyam, J. & Reuben, R. (1987) Survey of the anopheline fauna of the Western Ghats in Tamil Nadu, India. *Indian Journal of Malariology*, 24, 21–28.
- Theobald, F.V. (1903) *A Monograph of the Culicidae or Mosquitoes*, 3. British Museum (Natural History), London.
- Theobald, F.V. (1910) *A Monograph of the Culicidae or Mosquitoes*, 5. British Museum (Natural History), London.
- Tsukamoto, M., Miyata, A. & Miyagi, I. (1978) Surveys on simian malaria parasites and their vector in Palawan Island, the Philippines. *Tropical Medicine*, 20, 39–50.
- Tun-Lin, W., Aung, H., Moe, M., Sebastian, A., Paing, M. & Thu, M.-M. (1987) Some environmental factors influencing the breeding of *Anopheles balabacensis* complex (*dirus*) in domestic wells in Burma. *Journal of Communicable Diseases*, 19, 291–299.
- Tun-Lin, W., Thu, M.-M., Than, S.-M. & Mya, M.M. (1995) Hyperendemic malaria in a forested, hilly Myanmar village. *Journal of the American Mosquito Control Association*, 11, 401–407.
- Tun-Lin, W., Tin, A.-A., Pe, H. & Kanda, T. (1988) Allozyme polymorphism at the esterase loci of two taxa of the *Anopheles balabacensis* complex in Burma. *Mosquito Borne Disease Bulletin*, 4, 68–71.
- Walton, C., Handley, J.M., Collins, F.H., Baimai, V., Harbach, R.E., Deesin, V. & Butlin, R.K. (2001) Genetic population structure and introgression in *Anopheles dirus* mosquitoes in South-east Asia. *Molecular Ecology*, 10, 569–580.
- Walton, C., Handley, J.M., Kuvangkadilok, C., Collins, F.H., Harbach, R.E., Baimai, V. & Butlin, R.K. (1999) Identification of five species of the *Anopheles dirus* complex from Thailand, using allele-specific polymerase chain reaction. *Medical and Veterinary Entomology*, 13, 24–32.
- Walton, C., Handley, J.M., Tun-Lin, W., Collins, F.H., Harbach, R.E., Baimai, V. & Butlin, R.K. (2000) Population structure and population history of *Anopheles dirus* mosquitoes in Southeast Asia. *Molecular Biology and Evolution*, 17, 962–974.
- Ward, R.A. (1984) Second supplement to 'A catalog of the mosquitoes of the world.' (Diptera: Culicidae). *Mosquito Systematics*, 16, 227–270.
- Warren, McW. & Wharton, R.H. (1963) The vectors of simian malaria: identity, biology, and geographic distribution. *Journal of Parasitology*, 49, 892–904.
- White, G.B. (1983) The importance of *Anopheles leucosphyrus* group mosquitoes as vectors of malaria and filariasis in relation to transmigration and forestry in Indonesia with assessments of *Anopheles balabacensis* ecology and vectorial capacity. Assignment Report: 12 July – 15 September 1982. WHO Project IR VBC 025. World Health Organization, Vector Biology and Control Research Unit, Jakarta, Indonesia.
- Wibowo, S., Baimai, V. & Andre, R.G. (1984) Differentiation of four taxa of the *Anopheles balabacensis* complex using H-banding patterns in the sex chromosomes. *Canadian Journal of Genetics and Cytology*, 26, 425–429.
- Wilkerson, R.C. & Peyton, E.L. (1990) Standardized nomenclature for the costal wing spots of the genus *Anopheles* and other spotted-wing mosquitoes (Diptera: Culicidae). *Journal of Medical Entomology*, 27, 207–224.
- Xu, J.-N. & Qu, F.-Y. (1997) Ribosomal DNA difference between species A and D of the *Anopheles dirus* complex of mosquito from China. *Medical and Veterinary Entomology*, 11, 134–138.
- Xu, X.-C., Xu, J.-N. & Qu, F.-Y. (1998) A diagnostic polymerase chain reaction assay for species A and D of the *Anopheles dirus*

- (Diptera: Culicidae) species complex based on ribosomal DNA second internal transcribed spacer sequence. *Journal of the American Mosquito Control Association*, **14**, 385–389.
- Yasothornsrikul, S., Panyim, S. & Rosenberg, R. (1988) Diagnostic restriction fragment patterns of DNA from the four isomorphic species of *Anopheles dirus*. *Southeast Asian Journal of Tropical Medicine and Public Health*, **19**, 703–708.
- Yong, H.S., Cheong, W.H., Chiang, G.L., Dhaliwal, S.S., Loong, K.P. & Sarjan, K.P. (1983) Biochemical genetic relationship in three taxa of the malaria mosquito *Anopheles balabacensis* complex. *Comparative Biochemistry and Physiology B*, **76**, 611–613.
- Zahedi, M. & White, G.B. (1994) Filaria vector competence of some *Anopheles* species. *Tropical Medicine and Parasitology*, **45**, 27–32.
- Zulueta, J. de (1956) Malaria in Sarawak and Brunei. *Bulletin of the World Health Organization*, **15**, 65–671.
- Zulueta, J. de (1957) Observations on filariasis in Sarawak and Brunei. *Bulletin of the World Health Organization*, **16**, 699–705.
- Zulueta, J.A. de & LaChance, F. (1956) A malaria-control experiment in the interior of Borneo. *Bulletin of the World Health Organization*, **15**, 673–693.

Accepted 30 December 2004